

## Section 3.1 : Introduction to Determinants

Chapter 3 : Determinants

Math 1554 Linear Algebra

## Topics and Objectives

### Topics

We will cover these topics in this section.

1. The definition and computation of a determinant
2. The determinant of triangular matrices

### Objectives

For the topics covered in this section, students are expected to be able to do the following.

1. Compute determinants of  $n \times n$  matrices using a cofactor expansion.
2. Apply theorems to compute determinants of matrices that have particular structures.

Section 3.1 Slide 172

Section 3.1 Slide 173

## A Definition of the Determinant

Suppose  $A$  is  $n \times n$  and has elements  $a_{ij}$ .

1. If  $n = 1$ ,  $A = [a_{11}]$ , and has determinant  $\det A = a_{11}$ .
2. Inductive case: for  $n > 1$ ,

$$\det A = a_{11} \det A_{11} - a_{12} \det A_{12} + \cdots + (-1)^{1+n} a_{1n} \det A_{1n}$$

where  $A_{ij}$  is the submatrix obtained by eliminating row  $i$  and column  $j$  of  $A$ .

### Example

$$A = \begin{pmatrix} \bullet & \bullet & \bullet & \bullet & \bullet \\ \bullet & \bullet & \bullet & \bullet & \bullet \\ \bullet & \bullet & \bullet & \bullet & \bullet \\ \bullet & \bullet & \bullet & \bullet & \bullet \\ \bullet & \bullet & \bullet & \bullet & \bullet \end{pmatrix} \Rightarrow A_{2,3} = \begin{pmatrix} \bullet & \bullet & \bullet & \bullet \\ \bullet & \bullet & \bullet & \bullet \\ \bullet & \bullet & \bullet & \bullet \\ \bullet & \bullet & \bullet & \bullet \end{pmatrix}$$

Section 3.1 Slide 174

Section 3.1 Slide 175

## Example 1

$$\text{Compute } \det \begin{bmatrix} a & b \\ c & d \end{bmatrix}.$$

## Example 2

$$\text{Compute } \det \begin{bmatrix} 1 & -5 & 0 \\ 2 & 4 & -1 \\ 0 & 2 & 0 \end{bmatrix} = \begin{vmatrix} 1 & -5 & 0 \\ 2 & 4 & -1 \\ 0 & 2 & 0 \end{vmatrix}.$$

Section 3.1 Slide 176

### Theorem

The determinant of a matrix  $A$  can be computed down any row or column of the matrix. For instance, down the  $j^{\text{th}}$  column, the determinant is

$$\det A = a_{1j}C_{1j} + a_{2j}C_{2j} + \cdots + a_{nj}C_{nj}.$$

This gives us a way to calculate determinants more efficiently.

Section 3.1 Slide 178

## Cofactors

Cofactors give us a more convenient notation for determinants.

### Definition: Cofactor

The  $(i, j)$  cofactor of an  $n \times n$  matrix  $A$  is

$$C_{ij} = (-1)^{i+j} \det A_{ij}$$

The pattern for the negative signs is

$$\begin{pmatrix} + & - & + & - & \dots \\ - & + & - & + & \dots \\ + & - & + & - & \dots \\ - & + & - & + & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

Section 3.1 Slide 177

## Example 3

$$\text{Compute the determinant of } \begin{bmatrix} 5 & 4 & 3 & 2 \\ 0 & 1 & 2 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & 1 & 1 & 3 \end{bmatrix}.$$

Section 3.1 Slide 179



## Row Operations

- We saw how determinants are difficult or impossible to compute with a cofactor expansion for large  $N$ .
- Row operations give us a more efficient way to compute determinants.

### Theorem: Row Operations and the Determinant

Let  $A$  be a square matrix.

1. If a multiple of a row of  $A$  is added to another row to produce  $B$ , then  $\det B = \det A$ .
2. If two rows are interchanged to produce  $B$ , then  $\det B = -\det A$ .
3. If one row of  $A$  is multiplied by a scalar  $k$  to produce  $B$ , then  $\det B = k \det A$ .

**Example 1** Compute  $\begin{vmatrix} 1 & -4 & 2 \\ -2 & 8 & -9 \\ -1 & 7 & 0 \end{vmatrix}$

## Invertibility

Important practical implication: If  $A$  is reduced to echelon form, by  $r$  interchanges of rows and columns, then

$$|A| = \begin{cases} (-1)^r \times (\text{product of pivots}), & \text{when } A \text{ is invertible} \\ 0, & \text{when } A \text{ is singular.} \end{cases}$$

**Example 2** Compute the determinant

$$\begin{vmatrix} 0 & 1 & 2 & -1 \\ 2 & 5 & -7 & 3 \\ 0 & 3 & 6 & 2 \\ -2 & -5 & 4 & 2 \end{vmatrix}$$

## Properties of the Determinant

For any square matrices  $A$  and  $B$ , we can show the following.

1.  $\det A = \det A^T$ .
2.  $A$  is invertible if and only if  $\det A \neq 0$ .
3.  $\det(AB) = \det A \cdot \det B$ .

Section 3.2 Slide 188

## Additional Example (if time permits)

Use a determinant to find all values of  $\lambda$  such that matrix  $C$  is not invertible.

$$C = \begin{pmatrix} 5 & 0 & 0 \\ 0 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix} - \lambda I_3$$

Section 3.2 Slide 189

## Additional Example (if time permits)

Determine the value of

$$\det A = \det \left( \begin{pmatrix} 0 & 2 & 0 \\ 1 & 1 & 2 \\ 1 & 1 & 3 \end{pmatrix}^8 \right).$$

Section 3.2 Slide 190

## Section 3.3 : Volume, Linear Transformations

Chapter 3 : Determinants

Math 1554 Linear Algebra

Section 3.3 Slide 191

## Topics and Objectives

### Topics

We will cover these topics in this section.

1. Relationships between area, volume, determinants, and linear transformations.

### Objectives


For the topics covered in this section, students are expected to be able to do the following.

1. Use determinants to compute the area of a parallelogram, or the volume of a parallelepiped, possibly under a given linear transformation.

Students are not expected to be familiar with Cramer's rule.

## Determinants, Area and Volume

In  $\mathbb{R}^2$ , determinants give us the area of a parallelogram.



## Determinants as Area, or Volume

### Theorem

The volume of the parallelepiped spanned by the columns of an  $n \times n$  matrix  $A$  is  $|\det A|$ .

**Key Geometric Fact (which works in any dimension).** The area of the parallelogram spanned by two vectors  $\vec{a}, \vec{b}$  is equal to the area spanned by  $\vec{a}, c\vec{a} + \vec{b}$ , for any scalar  $c$ .




FIGURE 2 Two parallelograms of equal area.

Any  $3 \times 3$  matrix  $A$  can be transformed into a diagonal matrix using column operations that do not change  $|\det(A)|$ .




FIGURE 4 Two parallelepipeds of equal volume.

## Example 1

Calculate the area of the parallelogram determined by the points  $(-2, -2)$ ,  $(0, 3)$ ,  $(4, -1)$ ,  $(6, 4)$




FIGURE 5 Translating a parallelogram does not change its area.

## Linear Transformations

### Theorem

If  $T_A : \mathbb{R}^n \mapsto \mathbb{R}^n$ , and  $S$  is some parallelogram in  $\mathbb{R}^n$ , then

$$\text{volume}(T_A(S)) = |\det(A)| \cdot \text{volume}(S)$$

An example that applies this theorem is given in this week's worksheets.

## Section 4.9 : Applications to Markov Chains

Chapter 4 : Vector Spaces

Math 1554 Linear Algebra

## Topics and Objectives

### Topics

We will cover these topics in this section.

1. Markov chains
2. Steady-state vectors
3. Convergence

### Objectives


For the topics covered in this section, students are expected to be able to do the following.

1. Construct stochastic matrices and probability vectors.
2. Model and solve real-world problems using Markov chains (e.g. - find a steady-state vector for a Markov chain)
3. Determine whether a stochastic matrix is regular.

## Example 1

- A small town has two libraries,  $A$  and  $B$ .
- After 1 month, among the books checked out of  $A$ ,
  - 80% returned to  $A$
  - 20% returned to  $B$
- After 1 month, among the books checked out of  $B$ ,
  - 30% returned to  $A$
  - 70% returned to  $B$

If both libraries have 1000 books today, how many books does each library have after 1 month? After one year? After  $n$  months? A place to simulate this is <http://setosa.io/markov/index.html>



Section 4.9 Slide 200

## Example 1 Continued

The books are equally divided by between the two branches, denoted by  $\vec{x}_0 = \begin{bmatrix} .5 \\ .5 \end{bmatrix}$ . What is the distribution after 1 month, call it  $\vec{x}_1$ ? After two months?

After  $k$  months, the distribution is  $\vec{x}_k$ , which is what in terms of  $\vec{x}_0$ ?

Section 4.9 Slide 201

## Markov Chains

A few definitions:

- A **probability vector** is a vector,  $\vec{x}$ , with non-negative elements that sum to 1.
- A **stochastic matrix** is a square matrix,  $P$ , whose columns are probability vectors.
- A **Markov chain** is a sequence of probability vectors  $\vec{x}_k$ , and a stochastic matrix  $P$ , such that:

$$\vec{x}_{k+1} = P\vec{x}_k, \quad k = 0, 1, 2, \dots$$

- A **steady-state vector** for  $P$  is a vector  $\vec{q}$  such that  $P\vec{q} = \vec{q}$ .

Section 4.9 Slide 202

## Example 2

Determine a steady-state vector for the stochastic matrix

$$\begin{pmatrix} .8 & .3 \\ .2 & .7 \end{pmatrix}$$

Section 4.9 Slide 203



## Convergence

We often want to know what happens to a process,

$$\vec{x}_{k+1} = P\vec{x}_k, \quad k = 0, 1, 2, \dots$$

as  $k \rightarrow \infty$ .


**Definition:** a stochastic matrix  $P$  is **regular** if there is some  $k$  such that  $P^k$  only contains strictly positive entries.

### Theorem

If  $P$  is a regular stochastic matrix, then  $P$  has a unique steady-state vector  $\vec{q}$ , and  $\vec{x}_{k+1} = P\vec{x}_k$  converges to  $\vec{q}$  as  $k \rightarrow \infty$ .

## Stochastic Vectors in the Plane

The stochastic vectors in the plane are the line segment below, and a stochastic matrix maps stochastic vectors to themselves. Iterates  $P^k\vec{x}_0$  converge to the steady state.



$$P^k \vec{x}_0 \rightarrow [\vec{x}_\infty \quad \vec{x}_\infty]$$


## Example 3

A car rental company has 3 rental locations, A, B, and C. Cars can be returned at any location. The table below gives the pattern of rental and returns for a given week.

	rented from		
	A	B	C
returned to A	.8	.1	.2
returned to B	.2	.6	.3
returned to C	.0	.3	.5


There are 10 cars at each location today.

- Construct a stochastic matrix,  $P$ , for this problem.
- What happens to the distribution of cars after a long time? You may assume that  $P$  is regular.



$$P = \begin{bmatrix} .8 & .1 & .2 \\ .2 & .6 & .3 \\ .0 & .3 & .5 \end{bmatrix}$$

The Stochastic vectors in  $\mathbb{R}^3$ , are vectors  $\begin{bmatrix} s \\ t \\ 1-s-t \end{bmatrix}$ , where  $0 \leq s, t \leq 1$  and  $s+t \leq 1$ .  $P$  'contracts' stochastic vectors to  $x_\infty$ .



## Section 5.1 : Eigenvectors and Eigenvalues

Chapter 5 : Eigenvalues and Eigenvectors

Math 1554 Linear Algebra

### Topics and Objectives

#### Topics

We will cover these topics in this section.

1. Eigenvectors, eigenvalues, eigenspaces
2. Eigenvalue theorems

#### Objectives

For the topics covered in this section, students are expected to be able to do the following.

1. Verify that a given vector is an eigenvector of a matrix.
2. Verify that a scalar is an eigenvalue of a matrix.
3. Construct an eigenspace for a matrix.
4. Apply theorems related to eigenvalues (for example, to characterize the invertibility of a matrix).

### Eigenvectors and Eigenvalues

If  $A \in \mathbb{R}^{n \times n}$ , and there is a  $\vec{v} \neq \vec{0}$  in  $\mathbb{R}^n$ , and

$$A\vec{v} = \lambda\vec{v}$$

then  $\vec{v}$  is an **eigenvector** for  $A$ , and  $\lambda \in \mathbb{C}$  is the corresponding **eigenvalue**.

Note that

- We will only consider square matrices.
- If  $\lambda \in \mathbb{R}$ , then
  - when  $\lambda > 0$ ,  $A\vec{v}$  and  $\vec{v}$  point in the same direction
  - when  $\lambda < 0$ ,  $A\vec{v}$  and  $\vec{v}$  point in opposite directions
- Even when all entries of  $A$  and  $\vec{v}$  are real,  $\lambda$  can be complex (a rotation of the plane has no **real** eigenvalues.)
- We explore complex eigenvalues in Section 5.5.

## Example 1

Which of the following are eigenvectors of  $A = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$ ? What are the corresponding eigenvalues?

a)  $\vec{v}_1 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$

b)  $\vec{v}_2 = \begin{pmatrix} 1 \\ -1 \end{pmatrix}$

c)  $\vec{v}_3 = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$

Section 5.1 Slide 212

## Example 2

Confirm that  $\lambda = 3$  is an eigenvalue of  $A = \begin{pmatrix} 2 & -4 \\ -1 & -1 \end{pmatrix}$ .

Section 5.1 Slide 213

## Eigenspace

### Definition

Suppose  $A \in \mathbb{R}^{n \times n}$ . The eigenvectors for a given  $\lambda$  span a subspace of  $\mathbb{R}^n$  called the  $\lambda$ -**eigenspace** of  $A$ .

**Note:** the  $\lambda$ -eigenspace for matrix  $A$  is  $\text{Nul}(A - \lambda I)$ .

### Example 3

Construct a basis for the eigenspaces for the matrix whose eigenvalues are given, and sketch the eigenvectors.

$$\begin{pmatrix} 5 & -6 \\ 3 & -4 \end{pmatrix}, \quad \lambda = -1, 2$$

Section 5.1 Slide 214

## Theorems

Proofs for the following theorems are stated in Section 5.1. If time permits, we will explain or prove all/most of these theorems in lecture.

1. The diagonal elements of a triangular matrix are its eigenvalues.
2.  $A$  invertible  $\Leftrightarrow 0$  is not an eigenvalue of  $A$ .
3. If  $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_k$  are eigenvectors that correspond to distinct eigenvalues, then  $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_k$  are linearly independent.

Section 5.1 Slide 215

## Warning!

We can't determine the eigenvalues of a matrix from its reduced form.

Row reductions change the eigenvalues of a matrix.

**Example:** suppose  $A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$ . The eigenvalues are  $\lambda = 2, 0$ , because

$$A \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \end{bmatrix}$$
$$A \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

- But the reduced echelon form of  $A$  is:
- The reduced echelon form is triangular, and its eigenvalues are:

Section 5.1 Slide 216

## Additional Resource

### 3Blue1Brown

A beautiful, animated, and visual explanation of eigenvalues and eigenvectors.

<http://bit.ly/21XyJPg>

Section 5.1 Slide 217

## Section 5.2 : The Characteristic Equation

Chapter 5 : Eigenvalues and Eigenvectors

Math 1554 Linear Algebra

Section 5.2 Slide 218

## Topics and Objectives

### Topics

We will cover these topics in this section.

1. The characteristic polynomial of a matrix
2. Algebraic and geometric multiplicity of eigenvalues
3. Similar matrices

### Objectives

For the topics covered in this section, students are expected to be able to do the following.

1. Construct the characteristic polynomial of a matrix and use it to identify eigenvalues and their multiplicities.
2. Characterize the long-term behaviour of dynamical systems using eigenvalue decompositions.

Section 5.2 Slide 219

## The Characteristic Polynomial

### Recall:

$\lambda$  is an eigenvalue of  $A \Leftrightarrow (A - \lambda I)$  is not \_\_\_\_\_

Therefore, to calculate the eigenvalues of  $A$ , we can solve

$$\det(A - \lambda I) =$$

The quantity  $\det(A - \lambda I)$  is the **characteristic polynomial** of  $A$ .

The quantity  $\det(A - \lambda I) = 0$  is the **characteristic equation** of  $A$ .

The roots of the characteristic polynomial are the \_\_\_\_\_ of  $A$ .

## Example

The characteristic polynomial of  $A = \begin{pmatrix} 5 & 2 \\ 2 & 1 \end{pmatrix}$  is:

So the eigenvalues of  $A$  are:

## Characteristic Polynomial of $2 \times 2$ Matrices

Express the characteristic equation of

$$M = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

in terms of its determinant. What is the equation when  $M$  is singular?

## Algebraic Multiplicity

### Definition

The **algebraic multiplicity** of an eigenvalue is its multiplicity as a root of the characteristic polynomial.

### Example

Compute the algebraic multiplicities of the eigenvalues for the matrix

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

## Geometric Multiplicity

### Definition

The **geometric multiplicity** of an eigenvalue  $\lambda$  is the dimension of  $\text{Null}(A - \lambda I)$ .

1. Geometric multiplicity is always at least 1. It can be smaller than algebraic multiplicity.
2. Here is the basic example:

$$\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$$

$\lambda = 0$  is the only eigenvalue. Its algebraic multiplicity is 2, but the geometric multiplicity is 1.

## Example

Give an example of a  $4 \times 4$  matrix with  $\lambda = 0$  the only eigenvalue, but the geometric multiplicity of  $\lambda = 0$  is one.

## Recall: Long-Term Behavior of Markov Chains

### Recall:

- We often want to know what happens to a Markov Chain

$$\vec{x}_{k+1} = P\vec{x}_k, \quad k = 0, 1, 2, \dots$$

as  $k \rightarrow \infty$ .

- If  $P$  is regular, then there is a \_\_\_\_\_

### Now lets ask:

- If we don't know whether  $P$  is regular, what else might we do to describe the long-term behavior of the system?
- What can eigenvalues tell us about the behavior of these systems?


## Example: Eigenvalues and Markov Chains

*Note: the textbook has a similar example that you can review.*

Consider the Markov Chain:

$$\vec{x}_{k+1} = P\vec{x}_k = \begin{pmatrix} 0.6 & 0.4 \\ 0.4 & 0.6 \end{pmatrix} \vec{x}_k, \quad k = 0, 1, 2, 3, \dots, \quad \vec{x}_0 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

This system can be represented schematically with two nodes, A and B:



**Goal:** use eigenvalues to describe the long-term behavior of our system.

What are the eigenvalues of  $P$ ?

Use the eigenvalues and eigenvectors of  $P$  to analyze the long-term behaviour of the system. In other words, determine what  $\vec{x}_k$  tends to as  $k \rightarrow \infty$ .

What are the corresponding eigenvectors of  $P$ ?

## Similar Matrices

### Definition

Two  $n \times n$  matrices  $A$  and  $B$  are **similar** if there is a matrix  $P$  so that  $A = PBP^{-1}$ .

### Theorem

If  $A$  and  $B$  similar, then they have the same characteristic polynomial.

If time permits, we will explain or prove this theorem in lecture. Note:

- Our textbook introduces similar matrices in Section 5.2, but doesn't have exercises on this concept until 5.3.
- Two matrices,  $A$  and  $B$ , do not need to be similar to have the same eigenvalues. For example,

$$\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \text{ and } \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$$

## Additional Examples (if time permits)

1. True or false.
  - a) If  $A$  is similar to the identity matrix, then  $A$  is equal to the identity matrix.
  - b) A row replacement operation on a matrix does not change its eigenvalues.
2. For what values of  $k$  does the matrix have one real eigenvalue with algebraic multiplicity 2?

$$\begin{pmatrix} -3 & k \\ 2 & -6 \end{pmatrix}$$