

Sample Midterm 2B, Math 1554, Fall 2019

Instructors:

Date: Fall 2019, this is a 50 minute exam

PLEASE DO NOT PHOTOCOPY THIS EXAM

Section Number (e.g. A4, QH3, etc.) _____ TA Name _____

Student GT Email Address: _____@gatech.edu

First name (please write as legibly as possible within the boxes)														
Last name														
Student ID number														

Circle your instructor:

If you are in a QH section, which High School do you attend? _____

Student Instructions

- **Show your work** and justify your answers for all questions unless stated otherwise.
- **Organize your work** in a reasonably neat and coherent way.
- Calculators, notes, cell phones, books are not allowed.
- Use dark and clear writing: your exam will be scanned into a digital system.
- Exam pages are double sided. Be sure to complete both sides.
- Leave a 1 inch border around the edges of exams.
- Any work done on scratch paper will not be collected and will not be graded.


Math 1554, Sample Midterm 2B. Your initials: _____

You do not need to justify your reasoning for questions on this page.

1. (10 points) Indicate **true** if the statement is true, otherwise, indicate **false**.

	true	false
a) If $A\vec{x} = \vec{b}$ has exactly one solution for every \vec{b} , A must be singular.	<input type="radio"/>	<input type="radio"/>
b) If $\vec{x}, \vec{y} \in \mathbb{R}^3$ are linearly independent, then $\{\vec{x}, \vec{y}, \vec{x} + \vec{y}\}$ is a basis for \mathbb{R}^3 .	<input type="radio"/>	<input type="radio"/>
c) The set of solutions to $A\vec{x} = \vec{b}$, for any $\vec{b} \in \mathbb{R}^n$, is a subspace.	<input type="radio"/>	<input type="radio"/>
d) If $A, B \in \mathbb{R}^{n \times n}$ and $AB = I$, then $BA = I$.	<input type="radio"/>	<input type="radio"/>
e) Any matrix that is similar to the identity matrix must be equal to the identity matrix.	<input type="radio"/>	<input type="radio"/>
f) If $A, B \in \mathbb{R}^{m \times n}$ have the same null space, then they have the same RREF.	<input type="radio"/>	<input type="radio"/>
g) If A is $n \times n$, and there exists a $\vec{b} \in \mathbb{R}^n$ such that $A\vec{x} = \vec{b}$ is inconsistent, then $\det(A) = 0$.	<input type="radio"/>	<input type="radio"/>
h) If A has an LU factorization, then A is invertible.	<input type="radio"/>	<input type="radio"/>
i) If $A \in \mathbb{R}^{n \times n}$ has eigenvector \vec{x} then $2\vec{x}$ is also an eigenvector of A .	<input type="radio"/>	<input type="radio"/>
j) Swapping the rows of A does not change the value of $\det(A)$.	<input type="radio"/>	<input type="radio"/>

2. (2 points) A 2×2 matrix A has eigenvalues $\lambda_1 = 1$ and $\lambda_2 = 2$, with eigenvectors and eigenspaces indicated in the picture. Draw $A\vec{x}$ and $A\vec{y}$.



Math 1554, Sample Midterm 2B. Your initials: _____

You do not need to justify your reasoning for questions on this page.

3. (2 points) Fill in the missing entries of the 3×3 matrix A with **non-zero** numbers so that A has null space spanned by \vec{v} .

$$\vec{v} = \begin{pmatrix} 2 \\ -1 \\ -1 \end{pmatrix}, \quad A = \begin{pmatrix} 1 & 2 & 0 \\ 1 & 0 & 2 \\ 0 & - & - \end{pmatrix}$$

4. (6 points) If possible, write down an example of a matrix with the following properties. If it is not possible to do so, write *not possible*. You do not need to justify your reasoning.

- (a) A 4×3 matrix A with $\text{rank}(A) = 3$ and $\text{rank}(A^T) = 4$.

$$A = \begin{pmatrix} & & & & \\ & & & & \\ & & & & \\ & & & & \end{pmatrix}$$

- (b) A 2×3 matrix in RREF whose null space is spanned by $\begin{pmatrix} 2 \\ -4 \\ 1 \end{pmatrix}$.

$$A = \begin{pmatrix} & & & & \\ & & & & \end{pmatrix}$$

- (c) A 3×3 matrix in echelon form, A , such that $\text{Col}(A)$ is spanned by the vectors $\begin{pmatrix} 2 \\ 0 \\ 0 \end{pmatrix}$, $\begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix}$.

$$A = \begin{pmatrix} & & & & \\ & & & & \\ & & & & \end{pmatrix}$$

- (d) A 4×4 stochastic matrix, P , such that the Markov Chain $x_{k+1} = Px_k$ for $k = 0, 1, 2, \dots$, does not have a unique steady-state.

$$P = \begin{pmatrix} & & & & \\ & & & & \\ & & & & \\ & & & & \end{pmatrix}$$

5. (1 point) Suppose \vec{v}_1, \vec{v}_2 are eigenvectors of an 3×3 matrix A that correspond to eigenvalues λ_1 and λ_2 .

$$\vec{v}_1 = \begin{pmatrix} 1 \\ 8 \\ 0 \end{pmatrix}, \quad \vec{v}_2 = \begin{pmatrix} 5 \\ 0 \\ 2 \end{pmatrix}, \quad \lambda_1 = 1, \quad \lambda_2 = \frac{1}{10}$$

Vector \vec{p} is such that $\vec{p} = \vec{v}_1 - 13\vec{v}_2$. What does $A^k \vec{p}$ tend to as $k \rightarrow \infty$?

Math 1554, Sample Midterm 2B. Your initials: _____

You do not need to justify your reasoning for questions on this page.

6. (3 points) If the determinant $\left| \begin{pmatrix} a & b \\ 1 & 0 \end{pmatrix} \right| = 3$, compute the value of $\left| \begin{pmatrix} -1 & 0 & 0 \\ 2a & 2b & 0 \\ 0 & 0 & 5 \end{pmatrix} \right|$.

7. A is the 3×6 matrix $A = \begin{bmatrix} 1 & 6 & -4 & 0 & 0 & -1 \\ 0 & 0 & 0 & 1 & 2 & -6 \\ 0 & 0 & 0 & 0 & 1 & 6 \end{bmatrix}$

- (a) (1 point) The rank of A is _____.
- (b) (1 point) The dimension of $\text{Null}(A)$ is _____.
- (c) (2 points) Write down a basis for $\text{Col}(A)$.

- (d) (3 points) Construct a basis for $\text{Null}(A)$.

Math 1554, Sample Midterm 2B. Your initials: _____

8. (4 points) S is the parallelogram determined by $\vec{v}_1 = \begin{pmatrix} 4 \\ -2 \end{pmatrix}$, and $\vec{v}_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$. If $A = \begin{pmatrix} 2 & 3 \\ 2 & 2 \end{pmatrix}$, what is the area of the image of S under the map $\vec{x} \mapsto A\vec{x}$?

9. (4 points) If possible, compute the LU factorization of $A = \begin{pmatrix} 5 & 4 \\ 10 & 6 \\ 0 & 2 \\ -5 & 1 \end{pmatrix}$

Math 1554, Sample Midterm 2B. Your initials: _____

10. (4 points) List all possible values of k , if any, so that A has a real eigenvalue with geometric multiplicity 2. Show your work.

$$A = \begin{pmatrix} 3 & 2 & 0 \\ 2 & 3 & 0 \\ 0 & 0 & k \end{pmatrix}$$

11. (4 points) Construct a basis for the subspace

$$H = \{\vec{x} \in \mathbb{R}^3 : 5x_1 + 4x_2 - 7x_3 = 0\}.$$

Math 1554, Sample Midterm 2B. Your initials: _____

12. (5 points) A has only two distinct eigenvalues, 0 and 1. $A = \begin{pmatrix} 0 & 1 & -1 \\ 1 & 0 & 1 \\ 1 & -1 & 2 \end{pmatrix}$.

(a) Construct the eigenbasis for eigenvalue $\lambda = 0$.

(b) Construct the eigenbasis for eigenvalue $\lambda = 1$.