


LINEAR

ALGEBRA

Week






Midterm 2 Lecture Review Activity, Math 1554

Course Schedule

Cancelled lectures due to inclement weather will likely result in cancelling review lectures and possibly moving through

Week	Dates	Mon Lecture	Tue Studio	Wed Lecture	Thu Studio	Fri Lecture
1	8/21 - 8/25	1.1	WS1.1	1.2	WS1.2	1.3
2	8/28 - 9/1	1.4	WS1.3,1.4	1.5	WS1.5	1.7
3	9/4 - 9/8	Break	WS1.7	1.8	WS1.8	1.9
4	9/11 - 9/15	2.1	WS1.9,2.1	Exam 1 Review	Cancelled	2.2
5	9/18 - 9/22	2.3,2.4	WS2.2,2.3	2.5	WS2.4,2.5	2.8
6	9/25 - 9/29	2.9	WS2.8,2.9	3.1,3.2	WS3.1,3.2	3.3
7	10/2 - 10/6	4.9	WS3.3,4.9	5.1,5.2	WS5.1,5.2	5.2
8	10/9 - 10/13	Break	Break	Exam 2 Review	Cancelled	5.3
9	10/16 - 10/20	5.3	WS5.3	5.5	WS5.5	6.1
10	10/23 - 10/27	6.1,6.2	WS6.1	6.2	WS6.2	6.3
11	10/30 - 11/3	6.4	WS6.3,6.4	6.6,6.5	WS6.4,6.5	6.5
12	11/6 - 11/10	6.6	WS6.5,6.6	Exam 3 Review	Cancelled	PageRank
13	11/13 - 11/17	7.1	WSPageRank	7.2	WS7.1,7.2	7.3
14	11/20 - 11/24	7.3,7.4	WS7.2,7.3	Break	Break	Break
15	11/27 - 12/1	7.4	WS7.3,7.4	7.4	WS7.4	7.4
16	12/4 - 12/8	Last lecture	Last Studio	Reading Period		
17	12/11 - 12/15	Final Exam: MATH 1554 Common Final Exam Tuesday, December 12th at 6pm				

1. (3 points) T_A is the linear transform $x \rightarrow Ax$, $A \in \mathbb{R}^{2 \times 2}$, that projects points in \mathbb{R}^2 onto the x_2 -axis. Sketch the nullspace of A , the range of the transform, and the column space of A . How are the range and column space related to each other?



2. Indicate **true** if the statement is true, otherwise, indicate **false**.

true false

- a) $S = \{\vec{x} \in \mathbb{R}^3 \mid x_1 = a, x_2 = 4a, x_3 = x_1x_2\}$ is a subspace for any $a \in \mathbb{R}$. true false
- b) If A is square and non-zero, and $A\vec{x} = A\vec{y}$ for some $\vec{x} \neq \vec{y}$, then $\det(A) \neq 0$. true false

3. If possible, write down an example of a matrix or quantity with the given properties. If it is not possible to do so, write *not possible*.

(a) A is 2×2 , $\text{Col}A$ is spanned by the vector $\begin{pmatrix} 2 \\ 3 \end{pmatrix}$ and $\dim(\text{Null}(A)) = 1$. $A = \begin{pmatrix} & \\ & \end{pmatrix}$

(b) A is 2×2 , $\text{Col}A$ is spanned by the vector $\begin{pmatrix} 2 \\ 3 \end{pmatrix}$ and $\dim(\text{Null}(A)) = 0$. $A = \begin{pmatrix} & \\ & \end{pmatrix}$

(c) A is in RREF and $T_A : \mathbb{R}^3 \rightarrow \mathbb{R}^3$. The vectors u and v are a basis for the range of T .

$$u = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, v = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, A = \begin{pmatrix} & & \\ & & \\ & & \end{pmatrix}$$

4. Indicate whether the situations are possible or impossible by filling in the appropriate circle.

possible impossible

4.i) Vectors \vec{u} and \vec{v} are eigenvectors of square matrix A , and $\vec{w} = \vec{u} + \vec{v}$ is also an eigenvector of A .

4.ii) $T_A = A\vec{x}$ is one-to-one, $\dim(\text{Col}(A)) = 4$, and $T_A : \mathbb{R}^3 \rightarrow \mathbb{R}^4$.

5. (2 points) Fill in the blanks.

(a) If A is a 6×4 matrix in RREF and $\text{rank}(A) = 4$, what is the rank of A^T ?

(b) $T_A = A\vec{x}$, where $A \in \mathbb{R}^{2 \times 2}$, is a linear transform that first rotates vectors in \mathbb{R}^2 clockwise by π radians about the origin, then scales their x -component by a factor of 3, then projects them onto the x_1 -axis. What is the value of $\det(A)$?

6. (3 points) A virus is spreading in a lake. Every week,

- 20% of the healthy fish get sick with the virus, while the other healthy fish remain healthy but could get sick at a later time.
- 10% of the sick fish recover and can no longer get sick from the virus, 80% of the sick fish remain sick, and 10% of the sick fish die.

Initially there are exactly 1000 fish in the lake.

- a) What is the stochastic matrix, P , for this situation? Is P regular?
- b) Write down any steady-state vector for the corresponding Markov-chain.

6. (3 points) A virus is spreading in a lake. Every week,

- 20% of the healthy fish get sick with the virus, while the other healthy fish remain healthy but could get sick at a later time.
- 10% of the sick fish recover and can no longer get sick from the virus, 80% of the sick fish remain sick, and 10% of the sick fish die.

Initially there are exactly 1000 fish in the lake.

- a) What is the stochastic matrix, P , for this situation? Is P regular?
- b) Write down any steady-state vector for the corresponding Markov-chain.

Section 5.3 : Diagonalization

Chapter 5 : Eigenvalues and Eigenvectors

Math 1554 Linear Algebra

Motivation: it can be useful to take large powers of matrices, for example A^k , for large k .

But: multiplying two $n \times n$ matrices requires roughly n^3 computations. Is there a more efficient way to compute A^k ?

Topics and Objectives

Topics

1. Diagonal, similar, and diagonalizable matrices
2. Diagonalizing matrices

Learning Objectives

For the topics covered in this section, students are expected to be able to do the following.

1. Determine whether a matrix can be diagonalized, and if possible diagonalize a square matrix.
2. Apply diagonalization to compute matrix powers.

Section 5.3 : Diagonalization

Chapter 5 : Eigenvalues and Eigenvectors

Math 1554 Linear Algebra

Motivation: it can be useful to take large powers of matrices, for example A^k , for large k .

But: multiplying two $n \times n$ matrices requires roughly n^3 computations. Is there a more efficient way to compute A^k ?

Topics and Objectives

Topics

1. Diagonal, similar, and diagonalizable matrices
2. Diagonalizing matrices

Learning Objectives

For the topics covered in this section, students are expected to be able to do the following.

1. Determine whether a matrix can be diagonalized, and if possible diagonalize a square matrix.
2. Apply diagonalization to compute matrix powers.

Course Schedule

Cancellations due to inclement weather will likely result in cancelling review lectures and possibly moving through course material more quickly.

Week	Dates	Mon Lecture	Tue Studio	Wed Lecture	Thu Studio	Fri Lecture	
1	8/21 - 8/25	1.1	WS1.1	1.2	WS1.2	1.3	
2	8/28 - 9/1	1.4	WS1.3,1.4	1.5	WS1.5	1.7	
3	9/4 - 9/8	Break	WS1.7	1.8	WS1.8	1.9	
4	9/11 - 9/15	2.1	WS1.9,2.1	Exam 1, Review	Cancelled	2.2	
5	9/18 - 9/22	2.3,2.4	WS2.2,2.3	2.5	WS2.4,2.5	2.8	
6	9/25 - 9/29	2.9	WS2.8,2.9	3.1,3.2	WS3.1,3.2	3.3	
7	10/2 - 10/6	4.9	WS3.3,4.9	5.1,5.2	WS5.1,5.2	5.2	
8	10/9 - 10/13	Break	Break	Exam 2, Review	Cancelled	5.3	
9	10/16 - 10/20	5.3	WS5.3	5.5	WS5.5	6.1	
10	10/23 - 10/27	6.1,6.2	WS6.1	6.2	WS6.2	6.3	
11	10/30 - 11/3	6.4	WS6.3,6.4	6.6,6.5	WS6.6,6.5	6.5	
12	11/6 - 11/10	6.6	WS6.5,6.6	Exam 3, Review	Cancelled	PageRank	
13	11/13 - 11/17	7.1	WS7,PageRank	7.2	WS7.1,7.2	7.3	
14	11/20 - 11/24	7.3,7.4	WS7.2,7.3	Break	Break	Break	
15	11/27 - 12/1	7.4	WS7.3,7.4	7.4	WS7.4	7.4	
16	12/4 - 12/8	Last lecture	Last Studio	Reading Period			
17	12/11 - 12/15	Final Exams: MATH 1554 Common Final Exam Tuesday, December 12th at 6pm					

Section 5.3 Slide 202

Section 5.3 Slide 203

Diagonal Matrices

A matrix is **diagonal** if the only non-zero elements, if any, are on the main diagonal.

The following are all diagonal matrices.

$$\begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}, [2], I_n, \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

We'll only be working with diagonal square matrices in this course.

Powers of Diagonal Matrices

If A is diagonal, then A^k is easy to compute. For example,

$$A = \begin{pmatrix} 3 & 0 \\ 0 & 0.5 \end{pmatrix}$$

$$A^2 =$$

$$A^k =$$

But what if A is not diagonal?

Section 5.3 Slide 204

Section 5.3 Slide 205

Diagonalization

Suppose $A \in \mathbb{R}^{n \times n}$. We say that A is **diagonalizable** if it is similar to a diagonal matrix, D . That is, we can write

$$A = PDP^{-1}$$

Diagonalization

Theorem

If A is diagonalizable $\Leftrightarrow A$ has n linearly independent eigenvectors.

Note: the symbol \Leftrightarrow means "if and only if".

Also note that $A = PDP^{-1}$ if and only if

$$A = [\vec{v}_1 \ \vec{v}_2 \ \dots \ \vec{v}_n] \begin{bmatrix} \lambda_1 & & & \\ & \lambda_2 & & \\ & & \ddots & \\ & & & \lambda_n \end{bmatrix} [\vec{v}_1 \ \vec{v}_2 \ \dots \ \vec{v}_n]^{-1}$$

where $\vec{v}_1, \dots, \vec{v}_n$ are linearly independent eigenvectors, and $\lambda_1, \dots, \lambda_n$ are the corresponding eigenvalues (**in order**).

Distinct Eigenvalues

Theorem

If A is $n \times n$ and has n distinct eigenvalues, then A is diagonalizable.

Why does this theorem hold?

Is it necessary for an $n \times n$ matrix to have n distinct eigenvalues for it to be diagonalizable?

Non-Distinct Eigenvalues

Theorem. Suppose

- A is $n \times n$
- A has distinct eigenvalues $\lambda_1, \dots, \lambda_k$, $k \leq n$
- $a_i =$ algebraic multiplicity of λ_i
- $d_i =$ dimension of λ_i eigenspace ("geometric multiplicity")

Then

1. $d_i \leq a_i$ for all i
2. A is diagonalizable $\Leftrightarrow \sum d_i = n \Leftrightarrow d_i = a_i$ for all i
3. A is diagonalizable \Leftrightarrow the eigenvectors, for all eigenvalues, together form a basis for \mathbb{R}^n .

Example 1

Diagonalize if possible.

$$\begin{pmatrix} 2 & 6 \\ 0 & -1 \end{pmatrix}$$

Example 2

Diagonalize if possible.

$$\begin{pmatrix} 3 & 1 \\ 0 & 3 \end{pmatrix}$$

Example 3

The eigenvalues of A are $\lambda = 3, 1$. If possible, construct P and D such that $AP = PD$.

$$A = \begin{pmatrix} 7 & 4 & 16 \\ 2 & 5 & 8 \\ -2 & -2 & -5 \end{pmatrix}$$

Additional Example (if time permits)

Note that

$$\vec{x}_k = \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix} \vec{x}_{k-1}, \quad \vec{x}_0 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \quad k = 1, 2, 3, \dots$$

generates a well-known sequence of numbers.

Use a diagonalization to find a matrix equation that gives the n^{th} number in this sequence.

THEOREM 5**The Diagonalization Theorem**

An $n \times n$ matrix A is diagonalizable if and only if A has n linearly independent eigenvectors.

In fact, $A = PDP^{-1}$, with D a diagonal matrix, if and only if the columns of P are n linearly independent eigenvectors of A . In this case, the diagonal entries of D are eigenvalues of A that correspond, respectively, to the eigenvectors in P .

EXAMPLE 4 Diagonalize the following matrix, if possible.

$$A = \begin{bmatrix} 2 & 4 & 3 \\ -4 & -6 & -3 \\ 3 & 3 & 1 \end{bmatrix}$$

THEOREM 6

An $n \times n$ matrix with n distinct eigenvalues is diagonalizable.

THEOREM 7

Let A be an $n \times n$ matrix whose distinct eigenvalues are $\lambda_1, \dots, \lambda_p$.

- For $1 \leq k \leq p$, the dimension of the eigenspace for λ_k is less than or equal to the multiplicity of the eigenvalue λ_k .
- The matrix A is diagonalizable if and only if the sum of the dimensions of the eigenspaces equals n , and this happens if and only if (i) the characteristic polynomial factors completely into linear factors and (ii) the dimension of the eigenspace for each λ_k equals the multiplicity of λ_k .
- If A is diagonalizable and B_k is a basis for the eigenspace corresponding to λ_k for each k , then the total collection of vectors in the sets B_1, \dots, B_p forms an eigenvector basis for \mathbb{R}^n .

5.3 EXERCISES

In Exercises 1 and 2, let $A = PDP^{-1}$ and compute A^4 .

$$1. P = \begin{bmatrix} 5 & 7 \\ 2 & 3 \end{bmatrix}, D = \begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix}$$

$$2. P = \begin{bmatrix} 2 & -3 \\ -3 & 5 \end{bmatrix}, D = \begin{bmatrix} 1 & 0 \\ 0 & 1/2 \end{bmatrix}$$

In Exercises 3 and 4, use the factorization $A = PDP^{-1}$ to compute A^k , where k represents an arbitrary positive integer.

$$3. \begin{bmatrix} a & 0 \\ 3(a-b) & b \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 3 & 1 \end{bmatrix} \begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix} \begin{bmatrix} 1 & 0 \\ -3 & 1 \end{bmatrix}$$

$$4. \begin{bmatrix} -2 & 12 \\ -1 & 5 \end{bmatrix} = \begin{bmatrix} 3 & 4 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} -1 & 4 \\ 1 & -3 \end{bmatrix}$$

In Exercises 5 and 6, the matrix A is factored in the form PDP^{-1} . Use the Diagonalization Theorem to find the eigenvalues of A and a basis for each eigenspace.

$$5. \begin{bmatrix} 2 & 2 & 1 \\ 1 & 3 & 1 \\ 1 & 2 & 2 \end{bmatrix} =$$

$$\begin{bmatrix} 1 & 1 & 2 \\ 1 & 0 & -1 \\ 1 & -1 & 0 \end{bmatrix} \begin{bmatrix} 5 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1/4 & 1/2 & 1/4 \\ 1/4 & 1/2 & -3/4 \\ 1/4 & -1/2 & 1/4 \end{bmatrix}$$

$$6. \begin{bmatrix} 4 & 0 & -2 \\ 2 & 5 & 4 \\ 0 & 0 & 5 \end{bmatrix} =$$

$$\begin{bmatrix} -2 & 0 & -1 \\ 0 & 1 & 2 \\ 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} 5 & 0 & 0 \\ 0 & 5 & 0 \\ 0 & 0 & 4 \end{bmatrix} \begin{bmatrix} 0 & 0 & 1 \\ 2 & 1 & 4 \\ -1 & 0 & -2 \end{bmatrix}$$

Diagonalize the matrices in Exercises 7–20, if possible. The eigenvalues for Exercises 11–16 are as follows: (11) $\lambda = 1, 2, 3$; (12) $\lambda = 2, 8$; (13) $\lambda = 5, 1$; (14) $\lambda = 5, 4$; (15) $\lambda = 3, 1$; (16) $\lambda = 2, 1$. For Exercise 18, one eigenvalue is $\lambda = 5$ and one eigenvector is $(-2, 1, 2)$.

$$7. \begin{bmatrix} 1 & 0 \\ 6 & -1 \end{bmatrix}$$

$$8. \begin{bmatrix} 5 & 1 \\ 0 & 5 \end{bmatrix}$$

$$9. \begin{bmatrix} 3 & -1 \\ 1 & 5 \end{bmatrix}$$

$$10. \begin{bmatrix} 2 & 3 \\ 4 & 1 \end{bmatrix}$$

$$11. \begin{bmatrix} -1 & 4 & -2 \\ -3 & 4 & 0 \\ -3 & 1 & 3 \end{bmatrix}$$

$$12. \begin{bmatrix} 4 & 2 & 2 \\ 2 & 4 & 2 \\ 2 & 2 & 4 \end{bmatrix}$$

$$13. \begin{bmatrix} 2 & 2 & -1 \\ 1 & 3 & -1 \\ -1 & -2 & 2 \end{bmatrix}$$

$$14. \begin{bmatrix} 4 & 0 & -2 \\ 2 & 5 & 4 \\ 0 & 0 & 5 \end{bmatrix}$$

$$15. \begin{bmatrix} 7 & 4 & 16 \\ 2 & 5 & 8 \\ -2 & -2 & -5 \end{bmatrix}$$

$$16. \begin{bmatrix} 0 & -4 & -6 \\ -1 & 0 & -3 \\ 1 & 2 & 5 \end{bmatrix}$$

$$17. \begin{bmatrix} 4 & 0 & 0 \\ 1 & 4 & 0 \\ 0 & 0 & 5 \end{bmatrix}$$

$$18. \begin{bmatrix} -7 & -16 & 4 \\ 6 & 13 & -2 \\ 12 & 16 & 1 \end{bmatrix}$$

$$19. \begin{bmatrix} 5 & -3 & 0 & 9 \\ 0 & 3 & 1 & -2 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 2 \end{bmatrix}$$

$$20. \begin{bmatrix} 4 & 0 & 0 & 0 \\ 0 & 4 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 1 & 0 & 0 & 2 \end{bmatrix}$$

In Exercises 21 and 22, A , B , P , and D are $n \times n$ matrices. Mark each statement True or False. Justify each answer. (Study Theorems 5 and 6 and the examples in this section carefully before you try these exercises.)

21. a. A is diagonalizable if $A = PDP^{-1}$ for some matrix D and some invertible matrix P .
 b. If \mathbb{R}^n has a basis of eigenvectors of A , then A is diagonalizable.
 c. A is diagonalizable if and only if A has n eigenvalues, counting multiplicities.
 d. If A is diagonalizable, then A is invertible.
22. a. A is diagonalizable if A has n eigenvectors.
 b. If A is diagonalizable, then A has n distinct eigenvalues.
 c. If $AP = PD$, with D diagonal, then the nonzero columns of P must be eigenvectors of A .
 d. If A is invertible, then A is diagonalizable.
23. A is a 5×5 matrix with two eigenvalues. One eigenspace is three-dimensional, and the other eigenspace is two-dimensional. Is A diagonalizable? Why?