



7.1 Exercises

Determine which of the matrices in Exercises 1–6 are symmetric.

1. $\begin{bmatrix} 3 & 5 \\ 5 & -7 \end{bmatrix}$

2. $\begin{bmatrix} 3 & -5 \\ -5 & 3 \end{bmatrix}$

3. $\begin{bmatrix} 2 & 3 \\ 2 & 4 \end{bmatrix}$

4. $\begin{bmatrix} 0 & 8 & 3 \\ 8 & 0 & -4 \\ 3 & 2 & 0 \end{bmatrix}$

5. $\begin{bmatrix} -6 & 2 & 0 \\ 2 & -6 & 2 \\ 0 & 2 & -6 \end{bmatrix}$

6. $\begin{bmatrix} 1 & 2 & 2 & 1 \\ 2 & 2 & 2 & 1 \\ 2 & 2 & 1 & 2 \end{bmatrix}$

Determine which of the matrices in Exercises 7–12 are orthogonal. If orthogonal, find the inverse.

7. $\begin{bmatrix} .6 & .8 \\ .8 & -.6 \end{bmatrix}$

8. $\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$

9. $\begin{bmatrix} -4/5 & 3/5 \\ 3/5 & 4/5 \end{bmatrix}$

10. $\begin{bmatrix} 1/3 & 2/3 & 2/3 \\ 2/3 & 1/3 & -2/3 \\ 2/3 & -2/3 & 1/3 \end{bmatrix}$

11. $\begin{bmatrix} 2/3 & 2/3 & 1/3 \\ 0 & 1/3 & -2/3 \\ 5/3 & -4/3 & -2/3 \end{bmatrix}$

12. $\begin{bmatrix} .5 & .5 & -.5 & -.5 \\ .5 & .5 & .5 & .5 \\ .5 & -.5 & -.5 & .5 \\ .5 & -.5 & .5 & -.5 \end{bmatrix}$

Orthogonally diagonalize the matrices in Exercises 13–22, giving an orthogonal matrix P and a diagonal matrix D . To save

you time, the eigenvalues in Exercises 17–22 are the following:

- (17) $-4, 4, 7$; (18) $-3, -6, 9$; (19) $-2, 7$; (20) $-3, 15$; (21) $1, 5$,
9; (22) $3, 5$.

13. $\begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$

14. $\begin{bmatrix} 1 & -5 \\ -5 & 1 \end{bmatrix}$

15. $\begin{bmatrix} 3 & 4 \\ 4 & 9 \end{bmatrix}$

16. $\begin{bmatrix} 6 & -2 \\ -2 & 9 \end{bmatrix}$

17. $\begin{bmatrix} 1 & 1 & 5 \\ 1 & 5 & 1 \\ 5 & 1 & 1 \end{bmatrix}$

18. $\begin{bmatrix} 1 & -6 & 4 \\ -6 & 2 & -2 \\ 4 & -2 & -3 \end{bmatrix}$

19. $\begin{bmatrix} 3 & -2 & 4 \\ -2 & 6 & 2 \\ 4 & 2 & 3 \end{bmatrix}$

20. $\begin{bmatrix} 5 & 8 & -4 \\ 8 & 5 & -4 \\ -4 & -4 & -1 \end{bmatrix}$

21. $\begin{bmatrix} 4 & 3 & 1 & 1 \\ 3 & 4 & 1 & 1 \\ 1 & 1 & 4 & 3 \\ 1 & 1 & 3 & 4 \end{bmatrix}$

22. $\begin{bmatrix} 4 & 0 & 1 & 0 \\ 0 & 4 & 0 & 1 \\ 1 & 0 & 4 & 0 \\ 0 & 1 & 0 & 4 \end{bmatrix}$

23. Let $A = \begin{bmatrix} 4 & -1 & -1 \\ -1 & 4 & -1 \\ -1 & -1 & 4 \end{bmatrix}$ and $v = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$. Verify that 5 is

an eigenvalue of A and v is an eigenvector. Then orthogonally diagonalize A .

24. Let $A = \begin{bmatrix} 2 & -1 & 1 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{bmatrix}$, $v_1 = \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$, and $v_2 = \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$. Verify that v_1 and v_2 are eigenvectors of A . Then orthogonally diagonalize A .

In Exercises 25–32, mark each statement True or False (T/F). Justify each answer.

25. (T/F) An $n \times n$ matrix that is orthogonally diagonalizable must be symmetric.

26. (T/F) There are symmetric matrices that are not orthogonally diagonalizable.

27. (T/F) An orthogonal matrix is orthogonally diagonalizable.

28. (T/F) If $B = PDP^{-T}$, where $P^T = P^{-1}$ and D is a diagonal matrix, then B is a symmetric matrix.

29. (T/F) For a nonzero v in \mathbb{R}^n , the matrix vv^T is called a projection matrix.

30. (T/F) If $A^T = A$ and if vectors u and v satisfy $u \cdot v = 3u$ and $Av = 4v$, then $u \cdot v = 0$.

31. (T/F) An $n \times n$ symmetric matrix has n distinct real eigenvalues.

32. (T/F) The dimension of an eigenspace of a symmetric matrix is sometimes less than the multiplicity of the corresponding eigenvalue.

33. Show that if A is an $n \times n$ symmetric matrix, then $(Ax) \cdot y = x \cdot (Ay)$ for all x, y in \mathbb{R}^n .

34. Suppose A is a symmetric $n \times n$ matrix and B is any $n \times m$ matrix. Show that $B^T AB$, $B^T B$, and BB^T are symmetric matrices.

35. Suppose A is invertible and orthogonally diagonalizable. Explain why A^{-1} is also orthogonally diagonalizable.

36. Suppose A and B are both orthogonally diagonalizable and $AB = BA$. Explain why AB is also orthogonally diagonalizable.

37. Let $A = PDP^{-1}$, where P is orthogonal and D is diagonal, and let λ be an eigenvalue of A of multiplicity k . Then λ appears k times on the diagonal of D . Explain why the dimension of the eigenspace for λ is k .

38. Suppose $A = PRP^{-1}$, where P is orthogonal and R is upper triangular. Show that if A is symmetric, then R is symmetric and hence is actually a diagonal matrix.

39. Construct a spectral decomposition of A from Example 2.

40. Construct a spectral decomposition of A from Example 3.

41. Let u be a unit vector in \mathbb{R}^n , and let $B = uu^T$.

- a. Given any x in \mathbb{R}^n , compute Bx and show that Bx is the orthogonal projection of x onto u , as described in Section 6.2.

- b. Show that B is a symmetric matrix and $B^2 = B$.

- c. Show that u is an eigenvector of B . What is the corresponding eigenvalue?

42. Let B be an $n \times n$ symmetric matrix such that $B^2 = B$. Any such matrix is called a **projection matrix** (or an **orthogonal projection matrix**). Given any y in \mathbb{R}^n , let $\hat{y} = By$ and $z = y - \hat{y}$.

- a. Show that z is orthogonal to \hat{y} .

- b. Let W be the column space of B . Show that y is the sum of a vector in W and a vector in W^\perp . Why does this prove that By is the orthogonal projection of y onto the column space of B ?

Orthogonally diagonalize the matrices in Exercises 43–46. To practice the methods of this section, do not use an eigenvector routine from your matrix program. Instead, use the program to find the eigenvalues, and, for each eigenvalue λ , find an orthonormal basis for $\text{Nul}(A - \lambda I)$, as in Examples 2 and 3.

43. $\begin{bmatrix} 6 & 2 & 9 & -6 \\ 2 & 6 & -6 & 9 \\ 9 & -6 & 6 & 2 \\ -6 & 9 & 2 & 6 \end{bmatrix}$

44. $\begin{bmatrix} .63 & -.18 & -.06 & -.04 \\ -.18 & .84 & -.04 & .12 \\ -.06 & .04 & .72 & -.12 \\ -.04 & .12 & -.12 & .66 \end{bmatrix}$

45. $\begin{bmatrix} .31 & .58 & .08 & .44 \\ .58 & -.56 & .44 & -.58 \\ .08 & .44 & .19 & -.08 \\ .44 & -.58 & -.08 & .31 \end{bmatrix}$

46. $\begin{bmatrix} 8 & 2 & 2 & -6 & 9 \\ 2 & 8 & 2 & -6 & 9 \\ 2 & 2 & 8 & -6 & 9 \\ -6 & -6 & -6 & 24 & 9 \\ 9 & 9 & 9 & 9 & -21 \end{bmatrix}$

7.2 EXERCISES

1. Compute the quadratic form $\mathbf{x}^T A \mathbf{x}$, when $A = \begin{bmatrix} 5 & 1/3 \\ 1/3 & 1 \end{bmatrix}$
and
a. $\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$ b. $\mathbf{x} = \begin{bmatrix} 6 \\ 1 \end{bmatrix}$ c. $\mathbf{x} = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$
2. Compute the quadratic form $\mathbf{x}^T A \mathbf{x}$, for $A = \begin{bmatrix} 3 & 2 & 0 \\ 2 & 2 & 1 \\ 0 & 1 & 0 \end{bmatrix}$
and
5. Find the matrix of the quadratic form. Assume \mathbf{x} is in \mathbb{R}^3 .
a. $3x_1^2 + 2x_2^2 - 5x_3^2 - 6x_1x_2 + 8x_1x_3 - 4x_2x_3$
b. $6x_1x_2 + 4x_1x_3 - 10x_2x_3$
6. Find the matrix of the quadratic form. Assume \mathbf{x} is in \mathbb{R}^3 .
a. $3x_1^2 - 2x_2^2 + 5x_3^2 + 4x_1x_2 - 6x_1x_3$
b. $4x_3^2 - 2x_1x_2 + 4x_2x_3$
7. Make a change of variable, $\mathbf{x} = P\mathbf{y}$, that transforms the quadratic form $x_1^2 + 10x_1x_2 + x_2^2$ into a quadratic form with no cross-product term. Give P and the new quadratic form.
8. Let A be the matrix of the quadratic form
$$9x_1^2 + 7x_2^2 + 11x_3^2 - 8x_1x_2 + 8x_1x_3$$
- It can be shown that the eigenvalues of A are 3, 9, and 15. Find an orthogonal matrix P such that the change of variable $\mathbf{x} = P\mathbf{y}$ transforms $\mathbf{x}^T A \mathbf{x}$ into a quadratic form with no cross-product term. Give P and the new quadratic form.
- Classify the quadratic forms in Exercises 9–18. Then make a change of variable, $\mathbf{x} = P\mathbf{y}$, that transforms the quadratic form into one with no cross-product term. Write the new quadratic form. Construct P using the methods of Section 7.1.
9. $4x_1^2 - 4x_1x_2 + 4x_2^2$ 10. $2x_1^2 + 6x_1x_2 - 6x_2^2$
 11. $2x_1^2 - 4x_1x_2 - x_2^2$ 12. $-x_1^2 - 2x_1x_2 - x_2^2$
 13. $x_1^2 - 6x_1x_2 + 9x_2^2$ 14. $3x_1^2 + 4x_1x_2$
 15. [M] $-3x_1^2 - 7x_2^2 - 10x_3^2 - 10x_4^2 + 4x_1x_2 + 4x_1x_3 + 4x_1x_4 + 6x_3x_4$
 16. [M] $4x_1^2 + 4x_2^2 + 4x_3^2 + 4x_4^2 + 8x_1x_2 + 8x_3x_4 - 6x_1x_4 + 6x_2x_3$
 17. [M] $11x_1^2 + 11x_2^2 + 11x_3^2 + 11x_4^2 + 16x_1x_2 - 12x_1x_4 + 12x_2x_3 + 16x_3x_4$
 18. [M] $2x_1^2 + 2x_2^2 - 6x_1x_2 - 6x_1x_3 - 6x_1x_4 - 6x_2x_3 - 6x_2x_4 - 2x_3x_4$
 19. What is the largest possible value of the quadratic form $5x_1^2 + 8x_2^2$ if $\mathbf{x} = (x_1, x_2)$ and $\mathbf{x}^T \mathbf{x} = 1$, that is, if $x_1^2 + x_2^2 = 1$? (Try some examples of \mathbf{x} .)
 20. What is the largest value of the quadratic form $5x_1^2 - 3x_2^2$ if $\mathbf{x}^T \mathbf{x} = 1$?
21. a. The matrix of a quadratic form is a symmetric matrix.
b. A quadratic form has no cross-product terms if and only if the matrix of the quadratic form is a diagonal matrix.
c. The principal axes of a quadratic form $\mathbf{x}^T A \mathbf{x}$ are eigenvectors of A .
22. a. The expression $\|\mathbf{x}\|^2$ is not a quadratic form.
b. If A is symmetric and P is an orthogonal matrix, then the change of variable $\mathbf{x} = P\mathbf{y}$ transforms $\mathbf{x}^T A \mathbf{x}$ into a quadratic form with no cross-product term.
c. If A is a 2×2 symmetric matrix, then the set of \mathbf{x} such that $\mathbf{x}^T A \mathbf{x} = c$ (for a constant c) corresponds to either a circle, an ellipse, or a hyperbola.
d. An indefinite quadratic form is neither positive semidefinite nor negative semidefinite.
e. If A is symmetric and the quadratic form $\mathbf{x}^T A \mathbf{x}$ has only negative values for $\mathbf{x} \neq \mathbf{0}$, then the eigenvalues of A are all positive.
- Exercises 23 and 24 show how to classify a quadratic form $Q(\mathbf{x}) = \mathbf{x}^T A \mathbf{x}$, when $A = \begin{bmatrix} a & b \\ b & d \end{bmatrix}$ and $\det A \neq 0$, without finding the eigenvalues of A .
23. If λ_1 and λ_2 are the eigenvalues of A , then the characteristic polynomial of A can be written in two ways: $\det(A - \lambda I)$ and $(\lambda - \lambda_1)(\lambda - \lambda_2)$. Use this fact to show that $\lambda_1 + \lambda_2 = a + d$ (the diagonal entries of A) and $\lambda_1\lambda_2 = \det A$.
24. Verify the following statements.
a. Q is positive definite if $\det A > 0$ and $a > 0$.
b. Q is negative definite if $\det A > 0$ and $a < 0$.
c. Q is indefinite if $\det A < 0$.
25. Show that if B is $m \times n$, then $B^T B$ is positive semidefinite; and if B is $n \times n$ and invertible, then $B^T B$ is positive definite.
26. Show that if an $n \times n$ matrix A is positive definite, then there exists a positive definite matrix B such that $A = B^T B$. [Hint: Write $A = PDP^T$, with $P^T = P^{-1}$. Produce a diagonal matrix C such that $D = C^T C$, and let $B = PCP^T$. Show that B works.]

In Exercises 21 and 22, matrices are $n \times n$ and vectors are in \mathbb{R}^n . Mark each statement True or False. Justify each answer.

$$\text{a. } \mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \quad \text{b. } \mathbf{x} = \begin{bmatrix} -2 \\ -1 \\ 5 \end{bmatrix} \quad \text{c. } \mathbf{x} = \begin{bmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \\ 1/\sqrt{2} \end{bmatrix}$$

3. Find the matrix of the quadratic form. Assume \mathbf{x} is in \mathbb{R}^2 .

$$\text{a. } 3x_1^2 - 4x_1x_2 + 5x_2^2 \quad \text{b. } 3x_1^2 + 2x_1x_2$$

4. Find the matrix of the quadratic form. Assume \mathbf{x} is in \mathbb{R}^2 .

$$\text{a. } 5x_1^2 + 16x_1x_2 - 5x_2^2 \quad \text{b. } 2x_1x_2$$

d. A positive definite quadratic form Q satisfies $Q(\mathbf{x}) > 0$ for all \mathbf{x} in \mathbb{R}^n .

e. If the eigenvalues of a symmetric matrix A are all positive, then the quadratic form $\mathbf{x}^T A \mathbf{x}$ is positive definite.

f. A Cholesky factorization of a symmetric matrix A has the form $A = R^T R$, for an upper triangular matrix R with positive diagonal entries.

22. a. The expression $\|\mathbf{x}\|^2$ is not a quadratic form.

b. If A is symmetric and P is an orthogonal matrix, then the change of variable $\mathbf{x} = P\mathbf{y}$ transforms $\mathbf{x}^T A \mathbf{x}$ into a quadratic form with no cross-product term.

c. If A is a 2×2 symmetric matrix, then the set of \mathbf{x} such that $\mathbf{x}^T A \mathbf{x} = c$ (for a constant c) corresponds to either a circle, an ellipse, or a hyperbola.

d. An indefinite quadratic form is neither positive semidefinite nor negative semidefinite.

e. If A is symmetric and the quadratic form $\mathbf{x}^T A \mathbf{x}$ has only negative values for $\mathbf{x} \neq \mathbf{0}$, then the eigenvalues of A are all positive.

Exercises 23 and 24 show how to classify a quadratic form $Q(\mathbf{x}) = \mathbf{x}^T A \mathbf{x}$, when $A = \begin{bmatrix} a & b \\ b & d \end{bmatrix}$ and $\det A \neq 0$, without finding the eigenvalues of A .

23. If λ_1 and λ_2 are the eigenvalues of A , then the characteristic polynomial of A can be written in two ways: $\det(A - \lambda I)$ and $(\lambda - \lambda_1)(\lambda - \lambda_2)$. Use this fact to show that $\lambda_1 + \lambda_2 = a + d$ (the diagonal entries of A) and $\lambda_1\lambda_2 = \det A$.

24. Verify the following statements.

a. Q is positive definite if $\det A > 0$ and $a > 0$.

b. Q is negative definite if $\det A > 0$ and $a < 0$.

c. Q is indefinite if $\det A < 0$.

25. Show that if B is $m \times n$, then $B^T B$ is positive semidefinite; and if B is $n \times n$ and invertible, then $B^T B$ is positive definite.

26. Show that if an $n \times n$ matrix A is positive definite, then there exists a positive definite matrix B such that $A = B^T B$. [Hint: Write $A = PDP^T$, with $P^T = P^{-1}$. Produce a diagonal matrix C such that $D = C^T C$, and let $B = PCP^T$. Show that B works.]

EXERCISES

27. Let A and B be symmetric $n \times n$ matrices whose eigenvalues are all positive. Show that the eigenvalues of $A + B$ are all positive. [Hint: Consider quadratic forms.]
28. Let A be an $n \times n$ invertible symmetric matrix. Show that if the quadratic form $\mathbf{x}^T A \mathbf{x}$ is positive definite, then so is the quadratic form $\mathbf{x}^T A^{-1} \mathbf{x}$. [Hint: Consider eigenvalues.]

7.3 EXERCISES

In Exercises 1 and 2, find the change of variable $\mathbf{x} = P\mathbf{y}$ that transforms the quadratic form $\mathbf{x}^T A \mathbf{x}$ into $\mathbf{y}^T D \mathbf{y}$ as shown.

1. $5x_1^2 + 6x_2^2 + 7x_3^2 + 4x_1x_2 - 4x_2x_3 = 9y_1^2 + 6y_2^2 + 3y_3^2$
2. $3x_1^2 + 3x_2^2 + 5x_3^2 + 6x_1x_2 + 2x_1x_3 + 2x_2x_3 = 7y_1^2 + 4y_2^2$

Hint: \mathbf{x} and \mathbf{y} must have the same number of coordinates, so the quadratic form shown here must have a coefficient of zero for y_3^2 .

In Exercises 3–6, find (a) the maximum value of $Q(\mathbf{x})$ subject to the constraint $\mathbf{x}^T \mathbf{x} = 1$, (b) a unit vector \mathbf{u} where this maximum is attained, and (c) the maximum of $Q(\mathbf{x})$ subject to the constraints $\mathbf{x}^T \mathbf{x} = 1$ and $\mathbf{x}^T \mathbf{u} = 0$.

3. $Q(\mathbf{x}) = 5x_1^2 + 6x_2^2 + 7x_3^2 + 4x_1x_2 - 4x_2x_3$
(See Exercise 1.)

416 CHAPTER 7 Symmetric Matrices and Quadratic Forms

4. $Q(\mathbf{x}) = 3x_1^2 + 3x_2^2 + 5x_3^2 + 6x_1x_2 + 2x_1x_3 + 2x_2x_3$ (See Exercise 2.)
5. $Q(\mathbf{x}) = x_1^2 + x_2^2 - 10x_1x_2$
6. $Q(\mathbf{x}) = 3x_1^2 + 9x_2^2 + 8x_1x_2$
7. Let $Q(\mathbf{x}) = -2x_1^2 - x_2^2 + 4x_1x_2 + 4x_2x_3$. Find a unit vector \mathbf{x} in \mathbb{R}^3 at which $Q(\mathbf{x})$ is maximized, subject to $\mathbf{x}^T \mathbf{x} = 1$. [Hint: The eigenvalues of the matrix of the quadratic form Q are 2, -1, and -4.]
8. Let $Q(\mathbf{x}) = 7x_1^2 + x_2^2 + 7x_3^2 - 8x_1x_2 - 4x_1x_3 - 8x_2x_3$. Find a unit vector \mathbf{x} in \mathbb{R}^3 at which $Q(\mathbf{x})$ is maximized, subject to $\mathbf{x}^T \mathbf{x} = 1$. [Hint: The eigenvalues of the matrix of the quadratic form Q are 9 and -3.]
9. Find the maximum value of $Q(\mathbf{x}) = 7x_1^2 + 3x_2^2 - 2x_1x_2$, subject to the constraint $x_1^2 + x_2^2 = 1$. (Do not go on to find a vector where the maximum is attained.)
10. Find the maximum value of $Q(\mathbf{x}) = -3x_1^2 + 5x_2^2 - 2x_1x_2$, subject to the constraint $x_1^2 + x_2^2 = 1$. (Do not go on to find a vector where the maximum is attained.)
11. Suppose \mathbf{x} is a unit eigenvector of a matrix A corresponding to an eigenvalue 3. What is the value of $\mathbf{x}^T A \mathbf{x}$?
12. Let λ be any eigenvalue of a symmetric matrix A . Justify the statement made in this section that $m \leq \lambda \leq M$, where m and M are defined as in (2). [Hint: Find an \mathbf{x} such that $\mathbf{x} = \mathbf{x}^T A \mathbf{x}$.]
13. Let A be an $n \times n$ symmetric matrix, let M and m denote the maximum and minimum values of the quadratic form $\mathbf{x}^T A \mathbf{x}$, where $\mathbf{x}^T \mathbf{x} = 1$, and denote corresponding unit eigenvectors by \mathbf{u}_1 and \mathbf{u}_n . The following calculations show that given any number t between M and m , there is a unit vector \mathbf{x} such that $t = \mathbf{x}^T A \mathbf{x}$. Verify that $t = (1 - \alpha)m + \alpha M$ for some number α between 0 and 1. Then let $\mathbf{x} = \sqrt{1 - \alpha}\mathbf{u}_n + \sqrt{\alpha}\mathbf{u}_1$, and show that $\mathbf{x}^T \mathbf{x} = 1$ and $\mathbf{x}^T A \mathbf{x} = t$.

[M] In Exercises 14–17, follow the instructions given for Exercises 3–6.

14. $3x_1x_2 + 5x_1x_3 + 7x_1x_4 + 7x_2x_3 + 5x_2x_4 + 3x_3x_4$
15. $4x_1^2 - 6x_1x_2 - 10x_1x_3 - 10x_1x_4 - 6x_2x_3 - 6x_2x_4 - 2x_3x_4$
16. $-6x_1^2 - 10x_2^2 - 13x_3^2 - 13x_4^2 - 4x_1x_2 - 4x_1x_3 - 4x_1x_4 + 6x_3x_4$
17. $x_1x_2 + 3x_1x_3 + 30x_1x_4 + 30x_2x_3 + 3x_2x_4 + x_3x_4$

Exam 1 Out of 50 ...	Exam 2 Out of 50	Exam 3 Out of 50
30	26	17

Grade Calculator								
Assignment		Grade (%)		Weight	Points			
1	Exam (Best)	60.0	0.2	12.00			Exam 1	30 / 50
2	Exam (Mid)	52.0	0.2	10.40			Exam 2	26 / 50
3	Exam (Lowest)	34.0	0.15	5.10			Exam 3	17 / 50
4	MQE Score	10.00	0.25	25.00				
5	Final Exam	50.66667	0.2	10.13			Final Exam	48.67 / 100
6	Final Grade:	62.63						
7	Final Letter Grade:	D					In-studio Quiz Total	9 / 45
8							MyLab Total	172 / 214
9							Total Exploration Points	18 / 42
10								
11							MQE Total	70
12							MQE Score	70 / 70
13							MQE additional	0
14								
15								
16							Final Exam 2% Bonus? (CIO5 incentive)	TRUE
17							MQE Bonus	0
18							Total Final Exam Bonus	2
19								
20								
21								
22								

Exam 1 Out of 50 ...	Exam 2 Out of 50	Exam 3 Out of 50
34	34	25

Grade Calculator					
Assignment	Grade (%)	Weight	Points		
Exam (Best)	68.0	0.2	13.60	Exam 1	34 / 50
Exam (Mid)	68.0	0.2	13.60	Exam 2	34 / 50
Exam (Lowest)	50.0	0.15	7.50	Exam 3	25 / 50
MQE Score	100.0	0.25	25.00	Final Exam	62.00 / 100
Final Exam	64	0.2	12.80		
Final Grade:	72.50				
Final Letter Grade:	C			In-studio Quiz Total	9 / 45
				MyLab Total	172 / 214
				Total Exploration Points	18 / 42
11				MQE Total	70
12				MQE Score	70 / 70
13				MQE additional	0
14					
15					
16				Final Exam 2% Bonus? (Q10s incentive)	TRUE
17				MQE Bonus	0
18				Total Final Exam Bonus	2
19					
20					
21					
22					

Exam 1 Out of 50 ...	Exam 2 Out of 50	Exam 3 Out of 50
42	43	28

Exam 1 Out of 50 ...	Exam 2 Out of 50	Exam 3 Out of 50
34	28	34.5

Exam 1 Out of 50 ...	Exam 2 Out of 50	Exam 3 Out of 50
43	34	34.5

Grade Calculator				Grade Calculator				
				Grade Calculator				
				Grade Calculator				
A	B	C	D	E	F	G	H	I
1	Assignment	Grade (%)	Weight	Points				
2	Exam (Best)	86.0	0.2	17.20		Exam 1	43 /50	
3	Exam (Mid)	69.0	0.2	13.80		Exam 2	34 /50	
4	Exam (Lowest)	68.0	0.15	10.20		Exam 3	34.5 /50	
5	MQE Score	100.0	0.25	25.00				
6	Final Exam	79.85833	0.2	15.97		Final Exam	74.33 /100	
7	Final Grade:	B2.17						
8	Final Letter Grade:	B				In-studio Quiz Total	45 /45	
9						MyLab Total	214 /214	
10						Total Exploration Points	42 /42	
11								
12						MQE Total	140.5	
13						MQE Score	70 /70	
14						MQE additional	70.5	
15								
16						Final Exam 2% Bonus? (CIOs incentive)	TRUE	
17						MQE Bonus	3.525	
18						Total Final Exam Bonus	5.525	
19								
20								
21								
22								

Exam 1 Out of 50 ...	Exam 2 Out of 50	Exam 3 Out of 50
44.5	45	41.5

Course Schedule

Cancellations due to inclement weather will likely result in cancelling review lectures and possibly moving through course material at a faster pace.

Section 7.4 : The Singular Value Decomposition

Chapter 7: Orthogonality and Least Squares

Math 1554 Linear Algebra

$$4. \quad \begin{bmatrix} 3 & 0 \\ 8 & 3 \end{bmatrix} = U \Sigma V^T$$

orthogonal
 scale w/
 orthonorm.
 cols.

~~meta~~

Steps to compute SVD of A:

*compute $A^T A$

*find eigenvalues of $A^T A$ call them σ_i^2

*find orthonormal eigenvectors of $A^T A$ call them v_i

*Compute $u_i = 1/\sigma_i v_i A v_i$

$A = U \Sigma V^T$

$U = [u_1 \ u_2 \ \dots \ u_m]$ $V = [v_1 \ v_2 \ \dots \ v_n]$ both orthogonal matrices
 And Σ is a diagonal matrix with diagonal entries σ_i

Week Dates	Mon Lecture	Tue Studio	Wed Lecture	Thu Studio	Fri Lecture
1 8/19 - 8/23	1.1	WS1.1	1.2	WS1.2	1.3
2 8/26 - 8/30	1.4	WS1.3, 1.4	1.5	WS1.5	1.7
3 9/2 - 9/6	Break	WS1.7	1.8	WS1.8	1.9
4 9/9 - 9/13	2.1	WS1.9, 2.1	Exam 1, Review	Cancelled	2.2
5 9/16 - 9/20	2.3, 2.4	WS2.2, 2.3	2.5	WS2.4, 2.5	2.8
6 9/23 - 9/27	2.9	WS2.8, 2.9	2.9, 3.1	WS3.1	3.2
7 9/30 - 10/4	3.3	WS3.3	4.9	WS4.9	5.1
8 10/7 - 10/11	5.2	WS5.1, 5.2	Exam 2, Review	Cancelled	5.3
9 10/14 - 10/18	Break	Break	5.3	WS5.3	5.5
10 10/21 - 10/25	6.1	WS5.5, 6.1	6.2	WS6.2	6.3
11 10/28 - 11/1	6.4	WS6.3, 6.4	6.5	WS6.5	6.6
12 11/4 - 11/8	6.6, Review	WS6.6	Exam 3, Review	Cancelled	PageRank
13 11/11 - 11/15	7.1	WSPageRank	7.2	WS7.1, 7.2	7.3
14 11/18 - 11/22	7.3	WS7.3	7.4	WS7.4	7.4
15 11/25 - 11/29	7.4, Review	WS7.4, Review	Break	Break	Break
16 12/2 - 12/6	Last lecture	Last Studio	Reading Period		
17 12/5 - 12/12	Final Exams: MATH 1554 Common Final Exam 6:8:50pm on Tuesday, Dec. 10th				

$$U_1 = \frac{1}{\sqrt{5}} A v_1 \quad \left\{ \begin{array}{l} U_1 \cdot U_2 = 0 \\ \|U_1\| = 1 \end{array} \right.$$

$$U_2 = \frac{1}{\sqrt{2}} A v_2 \quad \left\{ \begin{array}{l} U_1 \cdot U_2 = 0 \\ \|U_2\| = 1 \end{array} \right.$$

Step 1: Compute $A^T A = \begin{bmatrix} 3 & 8 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 3 & 0 \\ 8 & 3 \end{bmatrix} = \begin{bmatrix} 73 & 24 \\ 24 & 9 \end{bmatrix}$

use $A^T A$ for U_1, U_2

$$P(\lambda) = \det(A - \lambda I) = \lambda^2 - 82\lambda + 81$$

$$= (\lambda - 81)(\lambda - 1) = 0$$

use $A^T A$

$$\lambda_1 = 81 \quad \lambda_2 = 1 \quad \text{(note } \lambda \in \mathbb{R})$$

$$\sigma_1 = 9 \quad \sigma_2 = 1$$

Step 2: Find v_1, v_2 orthonormal eigenvectors of $A^T A$

$$\lambda_1 = 81 \quad A^T A - 81I_2 = \begin{bmatrix} -8 & 24 \\ 24 & -72 \end{bmatrix} \sim \begin{bmatrix} 1 & -3 \\ 0 & 0 \end{bmatrix} \quad \hat{x} = \begin{bmatrix} 3 \\ 1 \end{bmatrix}$$

$$\lambda_2 = 1 \quad A^T A - I_2 = \begin{bmatrix} 72 & 24 \\ 24 & 8 \end{bmatrix} \sim \begin{bmatrix} 1 & 13 \\ 0 & 0 \end{bmatrix} \quad \hat{x} = \begin{bmatrix} -1/3 \\ 1 \end{bmatrix} \xrightarrow{\text{scale}} \begin{bmatrix} -1 \\ 3 \end{bmatrix} \quad v_2 = \begin{bmatrix} -1/\sqrt{10} \\ 3/\sqrt{10} \end{bmatrix}$$

$$\langle v_1, v_2 \rangle = 0?$$

$$\begin{bmatrix} 3 & 0 \\ 8 & 3 \end{bmatrix}$$

~~NEED~~

2x

Steps to compute SVD of A:

*compute $A^T A$

*find eigenvalues of $A^T A$ call them σ_i^{-2}

*find orthonormal eigenvectors of $A^T A$ call them v_i

*Compute $u_i = \frac{1}{\sigma_i} A v_i$

$$A = U \Sigma V^T$$

$U = [u_1 \ u_2 \ \dots \ u_m]$ $V = [v_1 \ v_2 \ \dots \ v_n]$ both orthogonal matrices

And Σ is a diagonal matrix with diagonal entries σ_i

~~NEED~~

$$\boxed{\begin{aligned} U_1 &= \frac{1}{\sigma_1} A v_1 & U_1 \cdot U_2 &= 0 \\ U_2 &= \frac{1}{\sigma_2} A v_2 & \|U_1\| &= 1 \\ && A^T &= A \end{aligned}}$$

$$\underline{\text{Step 1: Compute } A^T A = \begin{bmatrix} 3 & 8 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 3 & 0 \\ 8 & 3 \end{bmatrix}} = \begin{bmatrix} 73 & 24 \\ 24 & 9 \end{bmatrix}$$

use
 $A^T A$

for
 v_1, v_2

$$P(\lambda) = \det(A - \lambda I) = \lambda^2 - 82\lambda + 81$$

$$= (\lambda - 81)(\lambda - 1) = 0$$

$$\checkmark \quad \boxed{\lambda_1 = 81 \quad \lambda_2 = 1} \quad \text{use } \lambda's \quad (\text{note } \lambda \neq 0)$$

$$\boxed{\sigma_1 = 9 \quad \sigma_2 = 1}$$

$$\begin{aligned} 73(9) - (24)^2 \\ = 657 - 576 \\ = 81 \end{aligned}$$

Step 2: Find v_1, v_2 orthonormal eigenvectors of $A^T A$.

$$\lambda_1 = 81 \quad A^T A - 81I_2 = \begin{bmatrix} -8 & 24 \\ 24 & -72 \end{bmatrix} \sim \begin{bmatrix} 1 & -3 \\ 0 & 0 \end{bmatrix} \quad \vec{x} = \begin{pmatrix} 3 \\ 1 \end{pmatrix}$$

$$v_1 = \begin{pmatrix} 3/\sqrt{10} \\ 1/\sqrt{10} \end{pmatrix} \quad \checkmark$$

$$\uparrow v_1 \cdot v_2 = 0 ?$$

$$\lambda_2 = 1 \quad A^T A - I_2 = \begin{bmatrix} 72 & 24 \\ 24 & 8 \end{bmatrix} \sim \begin{bmatrix} 1 & 1/3 \\ 0 & 0 \end{bmatrix} \quad \vec{x} = \begin{pmatrix} -1/3 \\ 1 \end{pmatrix} \quad \xrightarrow{\text{(r2)} \sim} \begin{pmatrix} -1 \\ 3 \end{pmatrix} \quad v_2 = \begin{pmatrix} -1/\sqrt{10} \\ 3/\sqrt{10} \end{pmatrix}$$

Step 3: Compute u_1, u_2

$$u_1 = \frac{1}{\sigma_1} A v_1 = \frac{1}{9} \begin{bmatrix} 3 & 0 \\ 8 & 3 \end{bmatrix} \begin{pmatrix} 3/\sqrt{10} \\ 1/\sqrt{10} \end{pmatrix} = \frac{1}{9} \begin{pmatrix} 9/\sqrt{10} \\ 24/\sqrt{10} + 3/\sqrt{10} \end{pmatrix} = \begin{pmatrix} 1/\sqrt{10} \\ 3/\sqrt{10} \end{pmatrix} = \tilde{u}_1$$

$$u_2 = \frac{1}{\sigma_2} A v_2 = \frac{1}{1} \begin{bmatrix} 3 & 0 \\ 8 & 3 \end{bmatrix} \begin{pmatrix} -1/\sqrt{10} \\ 3/\sqrt{10} \end{pmatrix} = \frac{1}{1} \begin{pmatrix} -3/\sqrt{10} \\ 1/\sqrt{10} \end{pmatrix} = \begin{pmatrix} -3/\sqrt{10} \\ 1/\sqrt{10} \end{pmatrix} = \tilde{u}_2$$

So we have found

$$A = U \Sigma V^T$$

$$A = \begin{pmatrix} U_1 & U_2 \end{pmatrix} \begin{pmatrix} \sigma_1 & \\ & \sigma_2 \end{pmatrix} \begin{pmatrix} V_1^T & \\ & V_2^T \end{pmatrix}$$

recall from $A = P D P^T$

$$A = \lambda_1 U_1 U_1^T + \lambda_2 U_2 U_2^T$$

$$A = \sigma_1 U_1 V_1^T + \sigma_2 U_2 V_2^T$$

spectral decomp of A :

$$A = \begin{pmatrix} 3 & 0 \\ 8 & 3 \end{pmatrix}$$

$\text{Col } A = \mathbb{R}^2$

$\text{Row } A = \mathbb{R}^2$

$$U = \begin{pmatrix} 1/\sqrt{10} & -3/\sqrt{10} \\ 3/\sqrt{10} & 1/\sqrt{10} \end{pmatrix}$$

$$V = \begin{pmatrix} 3/\sqrt{10} & -1/\sqrt{10} \\ 1/\sqrt{10} & 3/\sqrt{10} \end{pmatrix}$$

$\text{rank } A = 2$

$\{U_1, V_1\}$ basis
for $\text{Row } A$

$\{U_2, V_2\}$ basis
for $\text{Col } A$

$$\begin{pmatrix} 3 & 0 \\ 8 & 3 \end{pmatrix} = 9 \begin{pmatrix} 1/\sqrt{10} \\ 3/\sqrt{10} \end{pmatrix} \begin{pmatrix} 3/\sqrt{10} & 1/\sqrt{10} \\ -1/\sqrt{10} & 3/\sqrt{10} \end{pmatrix} + 1 \begin{pmatrix} -3/\sqrt{10} \\ 1/\sqrt{10} \end{pmatrix} \begin{pmatrix} -1/\sqrt{10} & 3/\sqrt{10} \end{pmatrix}$$

$$= 9 \begin{pmatrix} 3/\sqrt{10} & 1/\sqrt{10} \\ 9/\sqrt{10} & 3/\sqrt{10} \end{pmatrix} + 1 \begin{pmatrix} 3/\sqrt{10} & -9/\sqrt{10} \\ -1/\sqrt{10} & 3/\sqrt{10} \end{pmatrix}$$

$$= \begin{pmatrix} 3 & 0 \\ 8 & 3 \end{pmatrix}$$

$$\begin{pmatrix} 2.7 & .9 \\ 8.1 & 2.7 \end{pmatrix} \quad \text{row 2}$$

$$\left\| \begin{pmatrix} 2.7 \\ .9 \\ 8.1 \\ 2.7 \end{pmatrix} - \begin{pmatrix} 3 \\ 0 \\ 8 \\ 3 \end{pmatrix} \right\| \quad \text{small at pos 2}$$

$$9. \begin{bmatrix} 3 & -3 \\ 0 & 0 \\ 1 & 1 \end{bmatrix} \text{ Step 1: } A^T A = \begin{pmatrix} 3 & 0 & 1 \\ -3 & 0 & 1 \end{pmatrix} \begin{pmatrix} 3 & -3 \\ 0 & 0 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 10 & -8 \\ -8 & 10 \end{pmatrix}$$

Row $A \in \mathbb{R}^2$

$\text{Col } A \subseteq \mathbb{R}^3$ 2-dim'l.

Step 2: Find U_1, U_2

$$\det(A) = \lambda^2 - 20\lambda + 36 = (\lambda - 18)(\lambda - 2) = 0 \quad \text{part of SVD}$$

$$\lambda_1 = 18 \quad \lambda_2 = 2$$

$$U_1 = \sqrt{18} = 3\sqrt{2}, \quad U_2 = \sqrt{2}$$

$$\boxed{\sqrt{1} \geq \sqrt{2}}$$

$$\lambda = 18 \quad A^T A - 18I_2 = \begin{pmatrix} -8 & -8 \\ -8 & -8 \end{pmatrix} \sim \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix} \quad \vec{x} = \begin{pmatrix} -1 \\ 1 \end{pmatrix} \quad U_1 = \begin{pmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \end{pmatrix}, \quad U_2 = \begin{pmatrix} 1/\sqrt{2} \\ -1/\sqrt{2} \end{pmatrix}$$

pack a unit length U_2
st. $U_1 \cdot U_2 = 0$

$$\text{Step 3: } U_1 = \frac{1}{\sqrt{2}} A v_1 = \frac{1}{\sqrt{2}} \begin{pmatrix} 3 & -3 \\ 0 & 0 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} -1/\sqrt{2} \\ 1/\sqrt{2} \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} -6/\sqrt{2} \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} -1 \\ 0 \\ 0 \end{pmatrix}$$

$$\text{basic for Col 1} \rightarrow U_2 = \frac{1}{\sqrt{2}} A v_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} 3 & -3 \\ 0 & 0 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ 2/\sqrt{2} \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \quad \frac{-6}{\sqrt{2}} \cdot \frac{1}{\sqrt{2}} = \frac{-6}{2} = -3$$

$$A = U \Sigma V^T = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 3\sqrt{2} & 0 \\ 0 & \sqrt{2} \\ 0 & 0 \end{pmatrix} \begin{pmatrix} -1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & 1/\sqrt{2} \end{pmatrix}$$

$\text{basis for Col } A$ is $\left\{ \begin{pmatrix} -1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \right\}$

$U_3 \in \text{span}\{v_1, v_2\}^\perp$

$+ \text{ set last column of } U \text{ we need}$

$$U = (U_1 \ U_2 \ U_3) \text{ to be orthogonal matrix}$$

(square 3×3 w/ orthonormal cols)

$$A = U_1 U_1 V^T + U_2 U_2 V^T$$

$$\begin{pmatrix} 3 & -3 \\ 0 & 0 \\ 1 & 1 \end{pmatrix} = 3\sqrt{2} \begin{pmatrix} -1 \\ 0 \\ 0 \end{pmatrix} (-1/\sqrt{2}, 1/\sqrt{2}) + \sqrt{2} \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} (1/\sqrt{2}, 1/\sqrt{2})$$

$$U_3 \cdot U_3 = 0 \quad U_2 \cdot U_3 = 0 \quad \text{if } \|U_3\| = 1$$

rank 2
over \mathbb{R}^2

$$= 3\sqrt{2} \begin{pmatrix} 1/\sqrt{2} & -1/\sqrt{2} \\ 0 & 0 \\ 0 & 0 \end{pmatrix} + \sqrt{2} \begin{pmatrix} 0 & 0 \\ 0 & 0 \\ 1/\sqrt{2} & 1/\sqrt{2} \end{pmatrix}$$

Topics and Objectives

Topics

1. The Singular Value Decomposition (SVD) and some of its applications.

Learning Objectives

1. Compute the SVD for a rectangular matrix.
2. Apply the SVD to
 - estimate the rank and condition number of a matrix,
 - construct a basis for the four fundamental spaces of a matrix, and
 - construct a spectral decomposition of a matrix.

Section 7.4 Slide 385

Col A Null(A^T)

$$\left[\begin{matrix} U_1 & U_2 & U_3 \end{matrix} \right] \left[\begin{matrix} \sigma_1 & 0 \\ 0 & \sigma_2 \\ 0 & 0 \end{matrix} \right] \left[\begin{matrix} V_1^T \\ V_2^T \end{matrix} \right]$$


Σ \sum V^T

Example 1

The linear transform whose standard matrix is

$$A = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 2\sqrt{2} & 0 \\ 0 & \sqrt{2} \end{pmatrix} = \begin{pmatrix} 2 & -1 \\ 2 & 1 \end{pmatrix}$$

maps the unit circle in \mathbb{R}^2 to an ellipse, as shown below. Identify the unit vector \vec{z} in which $\|A\vec{z}\|$ is maximized and compute this length.



Section 7.4 Slide 386

$$\sigma_j \mathbf{Av}_j = \mathbf{u}_j$$

Singular Values

$$\mathbf{Av}_j = \sigma_j \mathbf{u}_j$$

The matrix $A^T A$ is always symmetric, with non-negative eigenvalues $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_r \geq 0$. Let $\{\vec{v}_1, \dots, \vec{v}_n\}$ be the associated orthonormal eigenvectors. Then

$$\|A\vec{v}_j\|^2 = (\mathbf{Av}_j) \cdot (\mathbf{Av}_j) = \mathbf{v}_j^T A^T A \mathbf{v}_j = \lambda_j \quad \mathbf{v}_j \cdot \mathbf{v}_j = \lambda_j \quad \|\mathbf{v}_j\|^2 = \lambda_j$$

If the A has rank r , then $\{\mathbf{Av}_1, \dots, \mathbf{Av}_r\}$ is an orthogonal basis for $\text{Col } A$. For $1 \leq j < k \leq r$:

$$0 = (\mathbf{Av}_j)^T \mathbf{Av}_k = \mathbf{v}_j^T A^T A \mathbf{v}_k = \lambda_k \quad \mathbf{v}_j^T \mathbf{v}_k = \lambda_k \quad \mathbf{v}_j \cdot \mathbf{v}_k = 0. \quad \mathbf{v}_j, \mathbf{v}_k \text{ orthogonal.}$$

Definition: $\sigma_1 = \sqrt{\lambda_1} \geq \sigma_2 = \sqrt{\lambda_2} \geq \dots \geq \sigma_n = \sqrt{\lambda_n}$ are the singular values of A .

Section 7.4 Slide 388

$$\|\mathbf{Av}_j\|^2 = \lambda_j \Rightarrow \|\mathbf{Av}_j\| = \sqrt{\lambda_j} = \sigma_j$$

$$\left\| \frac{1}{\sigma_j} \mathbf{Av}_j \right\| = 1.$$


The SVD

Theorem: Singular Value Decomposition

A $m \times n$ matrix with rank r and non-zero singular values $\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_r$ has a decomposition $U\Sigma V^T$ where

$$\Sigma = \begin{bmatrix} D & 0 \\ 0 & 0 \end{bmatrix}_{m \times n} = \begin{bmatrix} \sigma_1 & 0 & \dots & 0 \\ 0 & \sigma_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \sigma_r \\ 0 & 0 & \dots & 0 \end{bmatrix}$$

U is a $m \times m$ orthogonal matrix, and V is a $n \times n$ orthogonal matrix.



$$M = U \cdot \Sigma \cdot V^*$$

$$\begin{pmatrix} \text{comple} \rightarrow \text{(same P)} \\ \downarrow \\ A = P D P^T \end{pmatrix}$$

↑ Symmetric

Defn.
a SVD $m \times n$ matrix A is given by

$$A = U \Sigma V^T$$

Where Σ is $m \times n$ (same size as A)
and diagonal.

U is $m \times m$ orthogonal } solve w/
 V^T is $n \times n$ orthogonal } orthonormal
cols

Algorithm to find the SVD of A

Suppose A is $m \times n$ and has rank $r \leq n$.

1. Compute the squared singular values of $A^T A$, σ_i^2 , and construct Σ . *by theory*.

Find eigenvalues of $A^T A$, call them λ_i ($i=1, \dots, r$) and set $\Sigma_i = \sqrt{\lambda_i}$.

2. Compute the unit singular vectors of $A^T A$, \vec{v}_i , use them to form V .

$A^T A$ symmetric, so by spectral theorem $\text{alg} = \text{geo}$

\Rightarrow we can find orthonormal basis $\{\vec{v}_1, \dots, \vec{v}_n\}$ of \mathbb{R}^m

3. Compute an orthonormal basis for $\text{Col } A$ using consisting of eigenvectors of $A^T A$.

$$\vec{u}_i = \frac{1}{\sigma_i} A \vec{v}_i, \quad i = 1, 2, \dots, r$$

Extend the set $\{\vec{u}_i\}$ to form an orthonormal basis for \mathbb{R}^m , use the basis to form U .

For any $\Sigma_i \neq 0$, compute

$$U_i = \{A \vec{v}_i\} \quad (\text{automatically unit length}).$$

$$\left[U_1 \ U_2 \ \dots \ U_r \ \underbrace{| \vec{u}_{r+1} \ \dots \ \vec{u}_m} \right] \underbrace{\begin{pmatrix} \Sigma_1 & & 0 \\ & \ddots & 0 \\ 0 & & \Sigma_{r+1} & 0 \\ & & & \ddots & 0 \\ & & & & 0 \end{pmatrix}}_{\text{fill in}} \left[V_1 \ \dots \ V_m \right]$$

s.t. U has orthonormal cols.

Example 3: Construct the singular value decomposition of

$$A = \begin{bmatrix} 1 & -1 \\ -2 & 2 \\ 2 & -2 \end{bmatrix}.$$

(It has rank 1.)

Example 2: Write down the singular value decomposition for

$$\begin{bmatrix} 2 & 0 \\ 0 & -3 \\ 0 & 0 \\ 0 & 0 \end{bmatrix} =$$

rank $A = 2$

Section 7.4

Slide 391

Section 7.4

Slide 392

Section 7.4

Slide 393

$\{U, \Sigma, V^T\}$ orthonormal basis of $\text{Null}(A)^\perp = (\text{Col } A)^\perp$

Example 2: Write down the singular value decomposition for

$$A = \begin{pmatrix} 2 & 0 \\ 0 & -3 \\ 0 & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} 2 & 0 & 0 & 0 \\ 0 & -3 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix} V^T$$

$\{U_1, U_2\}$ orthonormal basis for $\text{Col } A$

$\{V_1, V_2\}$ orthonormal basis of $\text{Row } A = \text{Null } A^\perp$

Step 1: Find λ 's

$$ATA = \begin{pmatrix} 2 & 0 & 0 & 0 \\ 0 & -3 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} 2 & 0 \\ 0 & -3 \\ 0 & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 4 & 0 \\ 0 & 9 \\ 0 & 0 \\ 0 & 0 \end{pmatrix} \quad \lambda_1 = 9, \lambda_2 = 4$$

$$\sqrt{\lambda_1} = 3 \geq \sqrt{\lambda_2} = 2$$

Step 2: Find U_1, V_1

$$\lambda_1 = 9 \quad ATA - 9I_2 = \begin{pmatrix} -5 & 0 \\ 0 & 0 \end{pmatrix} \quad \hat{x}_1 = \begin{pmatrix} 0 \\ 1 \end{pmatrix} \quad \hat{v}_1 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

$$\lambda_2 = 4 \quad V_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

(for just e_2)

fill in V_2 so V_1, V_2 are
one orthonormal set
of vectors.

Step 3: Find U_2, V_2

$$U_1 = \frac{1}{3} \begin{pmatrix} 2 & 0 \\ 0 & -3 \\ 0 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \frac{1}{3} \begin{pmatrix} 0 \\ -3 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ -1 \\ 0 \\ 0 \end{pmatrix}$$

$$U = \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$U_2 = \frac{1}{2} \begin{pmatrix} 2 & 0 \\ 0 & -3 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \frac{1}{2} \begin{pmatrix} 2 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$$

Find Null A .

$$\begin{pmatrix} U_1^T \\ U_2^T \end{pmatrix} = \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

Example 3: Construct the singular value decomposition of

$$A = \begin{pmatrix} 1 & -1 \\ -2 & 2 \\ 2 & -2 \end{pmatrix}.$$

(It has rank 1.)

Example 3: Construct the singular value decomposition of

$$A = \begin{bmatrix} 1 & -1 \\ -2 & 2 \\ 2 & -2 \end{bmatrix}$$

(It has rank 1.)

* non-full rank example

* what about wide matrices?

* four subspaces

* spectral decomp.

Step 1: Compute $A^T A$ & λ 's.

$$A^T A = \begin{bmatrix} 1 & -2 & 2 \\ -1 & 2 & -2 \\ 2 & -2 & 2 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ -2 & 2 \\ 2 & -2 \end{bmatrix} = \begin{bmatrix} 9 & -9 \\ -9 & 9 \end{bmatrix}$$

$$p(\lambda) = \lambda^2 - 18\lambda + 0 = (\lambda - 18)\lambda = 0$$

Section 7.4 Slide 393

$$\begin{aligned} \lambda_1 &= 18 & \lambda_2 &= 0 \\ \sigma_1 &= 3\sqrt{2} & \sigma_2 &= 0 \end{aligned}$$

$$A = U \sum V^T$$

need \vec{u}_2, \vec{v}_2

$$= \begin{pmatrix} -1/\sqrt{3} & \cdot & \cdot \\ 2/\sqrt{3} & \cdot & \cdot \\ -2/\sqrt{3} & \cdot & \cdot \end{pmatrix} \begin{pmatrix} 3\sqrt{2} & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} -1/\sqrt{2} & \vec{v}_2 \\ \vec{v}_2 & \vec{v}_2 \end{pmatrix}$$

Step 2: Find \vec{v}_1, \vec{v}_2

$$\stackrel{\text{def}}{=} A^T A - \lambda_1 I = \begin{bmatrix} 9 & -9 \\ -9 & 9 \end{bmatrix} - \begin{bmatrix} 18 & 0 \\ 0 & 18 \end{bmatrix} = \begin{bmatrix} -9 & -9 \\ -9 & -9 \end{bmatrix} \sim \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix} \quad \vec{x} = \begin{pmatrix} -1 \\ 1 \end{pmatrix} \quad \vec{v}_1 = \begin{pmatrix} -1/\sqrt{2} \\ 1/\sqrt{2} \end{pmatrix}$$

$$\stackrel{\text{def}}{=} A^T A - 0I = \begin{bmatrix} 9 & -9 \\ -9 & 9 \end{bmatrix} \sim \begin{pmatrix} 1 & -1 \\ 0 & 0 \end{pmatrix} \quad \vec{x} = \begin{pmatrix} 1 \\ 1 \end{pmatrix} \quad \vec{v}_2 = \begin{pmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \end{pmatrix} \quad \vec{v}_1 \cdot \vec{v}_2 = 0$$

Step 3: Find U_i 's $U_i = \frac{1}{\sigma_i} A \vec{v}_i$ for any nonzero σ_i 's.

$$U_1 = \frac{1}{3\sqrt{2}} \begin{pmatrix} 1 & -1 \\ -2 & 2 \\ 2 & -2 \end{pmatrix} \begin{pmatrix} -1/\sqrt{2} \\ 1/\sqrt{2} \end{pmatrix} = \frac{1}{3\sqrt{2}} \begin{pmatrix} -2/\sqrt{2} \\ 4/\sqrt{2} \\ -4/\sqrt{2} \end{pmatrix} = \frac{1}{3\sqrt{2}} \cdot \frac{1}{\sqrt{2}} \begin{pmatrix} -2 \\ 4 \\ -4 \end{pmatrix}$$

$$\|U_1\| = \sqrt{\left(-\frac{1}{3}\right)^2 + \left(\frac{2}{3}\right)^2 + \left(-\frac{2}{3}\right)^2} = \sqrt{\frac{1}{9} + \frac{4}{9} + \frac{4}{9}} = \sqrt{\frac{9}{9}} = \sqrt{1} = 1$$

option 1

Need U_2, U_3 that satisfy

$$\begin{aligned} U_1 \cdot U_2 &= 0 \\ U_1 \cdot U_3 &= 0 \\ U_2 \cdot U_3 &= 0 \end{aligned}$$

$$\|U_2\| = \|U_3\| = 1?$$

$$\begin{bmatrix} -1/\sqrt{3} & 1 & 0 & 0 \\ 2/\sqrt{3} & 0 & 1 & 0 \\ -2/\sqrt{3} & 0 & 0 & 1 \end{bmatrix} \sim \begin{bmatrix} -1/\sqrt{3} & 1 & 0 & 0 \\ 0 & 1 & * & * \\ 0 & 0 & 1 & * \end{bmatrix}$$

↑ e₁ e₂ e₃

Row reduce to find U_2, U_3

To GS on $\{U_1, U_2, U_3\}$

Option 3.

$$\text{Step 1: } \begin{bmatrix} U^T \\ I \end{bmatrix} = \begin{bmatrix} -\frac{1}{3} & \frac{2}{3} & -\frac{2}{3} \\ \frac{2}{3} & \frac{1}{3} & \frac{1}{3} \\ \frac{1}{3} & -\frac{2}{3} & \frac{2}{3} \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & -2 & 2 \\ 1 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad \vec{x} = r \begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix} + s \begin{pmatrix} -2 \\ 1 \\ 0 \end{pmatrix}$$

Do GS to $\left\{ \begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -2 \\ 1 \\ 0 \end{pmatrix} \right\}$.

$$V_1 = X_1 = \begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix}$$

$$V_2 = X_2 - \frac{x_2 \cdot v_1}{v_1 \cdot v_1} v_1 = \begin{pmatrix} -2 \\ 0 \\ 1 \end{pmatrix} - \frac{\begin{pmatrix} -2 \\ 0 \\ 1 \end{pmatrix} \cdot \begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix}}{\begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix} \cdot \begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix}} \begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix} = \begin{pmatrix} -2 \\ 0 \\ 1 \end{pmatrix} - \frac{-4}{5} \begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix}$$

$$= \begin{pmatrix} -2 \\ 0 \\ 1 \end{pmatrix} + \begin{pmatrix} 8/5 \\ 4/5 \\ 0 \end{pmatrix}$$

$$V_1 = \begin{pmatrix} 2/\sqrt{5} \\ 1/\sqrt{5} \\ 0 \end{pmatrix}$$

$$= \begin{pmatrix} -2/\sqrt{5} \\ 4/\sqrt{5} \\ 1 \end{pmatrix} \sim \begin{pmatrix} -2 \\ 4 \\ 5 \end{pmatrix}$$

$$V_2 = \begin{pmatrix} -2/\sqrt{45} \\ 4/\sqrt{45} \\ 5/\sqrt{45} \end{pmatrix}$$

So

$$U = \boxed{\begin{bmatrix} -\frac{1}{3} & \frac{2}{\sqrt{5}} & -\frac{2}{\sqrt{45}} \\ \frac{2}{3} & \frac{1}{\sqrt{5}} & \frac{4}{\sqrt{45}} \\ \frac{1}{3} & 0 & \frac{5}{\sqrt{45}} \end{bmatrix}}$$

THEOREM

The Invertible Matrix Theorem (concluded)

Let A be an $n \times n$ matrix. Then the following statements are each equivalent to the statement that A is an invertible matrix.

- u. $(\text{Col } A)^\perp = \{\mathbf{0}\}$.
- v. $(\text{Nul } A)^\perp = \mathbb{R}^n$.
- w. $\text{Row } A = \mathbb{R}^n$.
- x. A has n nonzero singular values.



IMT DLC no more updates.



Main page
Contents
Current events
Recent changes
About Wikipedia
Contact us
Donate

Contributor
Help

Learn to edit
Community portal
Recent changes
Upload file

Tools

What links here
Related changes

Special pages

Permanent link

Page information

Cite this page

What links here

Print/export

Download as PDF

Printable version

Languages

Arabic

Deutsch

Español

Singular value decomposition

From Wikipedia, the free encyclopedia

In linear algebra, the singular value decomposition (SVD) is a factorization of a real or complex matrix. It generalizes the eigendecomposition of a square normal matrix with an orthonormal eigenbasis to any $m \times n$ matrix. It is related to the polar decomposition.

Specifically, the singular value decomposition of an $m \times n$ complex matrix M is a factorization of the form $U\Sigma V^*$, where U is an $m \times m$ complex unitary matrix, Σ is an $m \times n$ rectangular diagonal matrix with non-negative real numbers on the diagonal, and V is an $n \times n$ complex unitary matrix. If M is real, U and V can also be guaranteed to be real orthogonal matrices. In such cases, the SVD is often denoted $U\Sigma V^t$.

The diagonal entries $\sigma_i = \Sigma_{ii}$ of Σ are known as the singular values of M . The number of non-zero singular values is equal to the rank of M . The columns of U and the columns of V are called the left-singular vectors and right-singular vectors of M , respectively.

The SVD is not unique. It is always possible to choose the decomposition so that the singular values Σ_{ii} are in descending order. In this case, Σ (but not always U and V) is uniquely determined by M .

The term sometimes refers to the compact SVD, a similar decomposition $M = U\Sigma V^*$ in which Σ is square diagonal of size $r \times r$, where $r \leq \min(m, n)$ is the rank of M , and has only the non-zero singular values. In this variant, U is an $m \times r$ semi-unitary matrix and V is an $n \times r$ semi-unitary matrix, such that $U^*U = V^*V = I_r$.

Mathematical applications of the SVD include computing the pseudoinverse, matrix approximation, and determining the rank, range, and null space of a matrix. The SVD is also extremely useful in all areas of science, engineering, and statistics, such as signal processing, least squares fitting of data, and process control.




Illustration of the singular value decomposition (SVD) of a real 2x2 matrix M .

Top: The action of M , indicated by its effect on the unit disc D and the two corresponding singular vectors e_1 and e_2 .

Left: The action of V^* , a rotation, on e_1 and e_2 .

Bottom: The action of Σ , a scaling by the singular values σ_1 horizontally and σ_2 vertically.

Right: The action of U , another rotation.

Contents [hide]

- 1 Intuitive interpretations
- 1.1 Rotation, coordinate scaling, and reflection
- 1.2 Singular values as semiaxes of an ellipse or ellipsoid
- 1.3 The columns of U and V are orthonormal bases
- 1.4 Geometric meaning
- 2 Example
- 3 SVD and spectral decomposition
- 3.1 Singular values

Applications of the SVD

The SVD has been applied to many modern applications in CS, engineering, and mathematics (our textbook mentions the first four).

- Estimating the rank and condition number of a matrix
- Constructing bases for the four fundamental spaces computing the pseudoinverse of a matrix
- Linear least squares problems
- Non-linear least-squares
 - https://en.wikipedia.org/wiki/Non-linear_least_squares
- Machine learning and data mining
 - <https://en.wikipedia.org/wiki/K-SVD>
- Facial recognition
 - <https://en.wikipedia.org/wiki/Eigenface>
- Principle component analysis
 - https://en.wikipedia.org/wiki/Principal_component_analysis
- Image compression

Students are expected to be familiar with the 1st two items in the list.

The Condition Number of a Matrix

If A is an invertible $n \times n$ matrix, the ratio

$$\frac{\sigma_1}{\sigma_n}$$

is the condition number of A .

Note that:

- The condition number of a matrix describes the sensitivity of a solution to $Ax = b$ to errors in A .
- We could define the condition number for a rectangular matrix, but that would go beyond the scope of this course.

Example 4

For $A = USV^T$, determine the rank of A , and orthonormal bases for $\text{Null}(A)$ and $(\text{Col}(A))^\perp$.

$$U = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\Sigma = \begin{bmatrix} 3 & 0 & 0 & 0 \\ 0 & \sqrt{0.8} & 0 & 0 \\ 0 & 0 & \sqrt{0.2} & 0 \\ 0 & 0 & 0 & \sqrt{0.2} \end{bmatrix}$$

$$V^T = \begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & \sqrt{0.8} \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & \sqrt{0.2} \end{bmatrix}$$

$$A = U \Sigma V^T$$

$$\text{rank } A = \text{rank } \Sigma = 3$$

$C \neq \text{non-zero } \sigma_i^2$

The Four Fundamental Spaces




FIGURE 4 The four fundamental subspaces and the action of A .

The Spectral Decomposition of a Matrix

The SVD can also be used to construct the spectral decomposition for any matrix with rank r

$$A = \sum_{i=1}^r \sigma_i \tilde{u}_i \tilde{v}_i^T,$$

where \tilde{u}_i, \tilde{v}_i are the i^{th} columns of U and V respectively.

For the case when $A = A^T$, we obtain the same spectral decomposition that we encountered in Section 7.2.

Section 7.4 Slide 399

Section 7.4 Slide 400

orthonormal basis of Row A is

$$\left\{ \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \\ 0 \end{bmatrix} \right\}$$

$$U_1 \in (\text{Null}(A))^\perp = \text{Row } A.$$

$$\textcircled{1} \quad A = \begin{bmatrix} 3 & -3 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} 3 & 0 \\ 0 & \sqrt{2} \end{bmatrix} \begin{bmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & -1/\sqrt{2} \end{bmatrix}^T$$

$$\textcircled{2} \quad A = \begin{bmatrix} 3 & 0 \\ 0 & 3 \end{bmatrix} = \begin{bmatrix} 1/\sqrt{10} & -3/\sqrt{10} \\ 3/\sqrt{10} & 1/\sqrt{10} \end{bmatrix} \begin{bmatrix} 9 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 3/\sqrt{10} & -1/\sqrt{10} \\ 1/\sqrt{10} & 3/\sqrt{10} \end{bmatrix}^T$$

orthonormal basis of $(\text{Row } A)^\perp = \text{Null } A$

$$\left\{ \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ -5/8 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right\}$$

$$U_4 \in \text{Null } A^T A$$

$$A = \begin{bmatrix} -1/\sqrt{3} & 2/\sqrt{5} & -2/\sqrt{5} \\ 2/\sqrt{3} & \sqrt{5}/5 & 4/\sqrt{5} \\ -2/\sqrt{3} & 0 & 5/\sqrt{5} \end{bmatrix} \begin{bmatrix} 35/2 & 0 \\ 0 & 0 \\ 0 & 0 \end{bmatrix} \begin{pmatrix} -1/\sqrt{2} & \sqrt{1/2} \\ \sqrt{1/2} & \sqrt{1/2} \end{pmatrix}$$

$$\xrightarrow{U_4 = 0} \xrightarrow{\lambda_4 = 0} \text{Null } A^T A = \text{Null } A \checkmark$$

orthonormal basis of $\text{Col } A$ is

$$\left\{ \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \\ 0 \end{bmatrix} \right\}$$

orthonormal basis for $(\text{Col } A)^\perp = \text{Null } A^T$

$$\left\{ \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \\ 0 \end{bmatrix} \right\}$$

$$U_1 = \frac{1}{\sqrt{1}} A v_1 \in \text{Col } A \quad \checkmark$$

$\text{in Col } A.$

$$U_4 \in (\text{Col } A)^\perp \quad \checkmark$$

$$A = \begin{pmatrix} 3 & -3 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} 3 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} 1/2 & 1/2 \\ 1/2 & 1/2 \\ 1/2 & 1/2 \end{pmatrix}^T$$

Find the

SVD of "wide"
matrix B .

See!

Compute $B^T B$ is 3×3 .

$$B = \begin{pmatrix} 3 & 0 & 1 \\ -3 & 0 & 1 \end{pmatrix}$$

$$\begin{pmatrix} 3 & -3 \\ 0 & 0 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 3 & 0 & 1 \\ -3 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 18 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 2 \end{pmatrix}$$

$$B^T = A = \begin{pmatrix} 3 & -3 \\ 0 & 0 \end{pmatrix}$$

skip this

$$= \begin{pmatrix} * & * & * \\ * & * & * \\ * & * & * \end{pmatrix}$$

$A^T A$ small matrix 2×2 .

$$A^T = (U \Sigma V^T)^T = (V^T \underline{\Sigma^+ U^T})$$

$$= \boxed{V \Sigma^T U^T}$$

is an SVD
of A^T .

$$A = \begin{pmatrix} 3 & -3 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 3\sqrt{2} & 0 \\ 0 & \sqrt{2} \\ 0 & 0 \end{pmatrix} \begin{pmatrix} -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix}^T$$

some place
for $A^T = A$

$$A = P D P^T$$

Q. Complete the spectral decompose.

$$A = \lambda_1 U_1 V_1^T + \lambda_2 U_2 V_2^T$$

$$A = \sqrt{\lambda}_1 U_1 V_1^T + \sqrt{\lambda}_2 U_2 V_2^T$$




$$= 3\sqrt{2} \begin{pmatrix} -1 \\ 0 \\ 0 \end{pmatrix} \begin{pmatrix} -1/\sqrt{2} & 1/\sqrt{2} \end{pmatrix} + \sqrt{2} \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \begin{pmatrix} 1/\sqrt{2} & 1/\sqrt{2} \end{pmatrix}$$

$$= 3\sqrt{2} \begin{pmatrix} 1/\sqrt{2} & -1/\sqrt{2} \\ 0 & 0 \\ 0 & 0 \end{pmatrix} + \sqrt{2} \begin{pmatrix} 0 & 0 \\ 0 & 0 \\ 1/\sqrt{2} & 1/\sqrt{2} \end{pmatrix}$$

```

clc
RGB=imread('buzz.jpg');
gray=rgb2gray(RGB);
A=im2double(gray);
[U,S,V]=svd(A);
sz=size(A)
rank(A)
Approx=zeros(sz);
r=2
for i=1:r
    u=U(:,i);
    s=S(i,i);
    v=V(:,i);
    Approx=Approx+s*u*v';
end
Approx;
% subplot(1,2,1),imshow(A),title('original');
% subplot(1,2,2),imshow(Approx),title(['low rank r=',num2str(r)]);


```



```

clc
RGB=imread('buzz.jpg');
gray=rgb2gray(RGB);
A=im2double(gray);
[U,S,V]=svd(A);
sz=size(A)
rank(A)
Approx=zeros(sz);
r=2
for i=1:r
    u=U(:,i);
    s=S(i,i);
    v=V(:,i);
    Approx=Approx+s*u*v';
end
Approx;
% subplot(1,2,1),imshow(A,title('original'));
% subplot(1,2,2),imshow(Approx),title(['low rank r=',num2str(r)]);

```



$$A\vec{x} = \lambda \vec{x}$$

$$(?)$$


* A Square

* A real

+ λ real

+ no restriction
on A

$$A\vec{v} = \sigma \vec{u}$$



L



7.4 EXERCISES

Find the singular values of the matrices in Exercises 1–4.

1. $\begin{bmatrix} 1 & 0 \\ 0 & -3 \end{bmatrix}$

2. $\begin{bmatrix} -3 & 0 \\ 0 & 0 \end{bmatrix}$

3. $\begin{bmatrix} 2 & 3 \\ 0 & 2 \end{bmatrix}$

4. $\begin{bmatrix} 3 & 0 \\ 8 & 3 \end{bmatrix}$

Find an SVD of each matrix in Exercises 5–12. [Hint: In Exercise 11, one choice for U is $\begin{bmatrix} -1/3 & 2/3 & 2/3 \\ 2/3 & -1/3 & 2/3 \\ 2/3 & 2/3 & -1/3 \end{bmatrix}$. In Exercise 12, one column of U can be $\begin{bmatrix} 1/\sqrt{6} \\ -2/\sqrt{6} \\ 1/\sqrt{6} \end{bmatrix}$.]

5. $\begin{bmatrix} -2 & 0 \\ 0 & 0 \end{bmatrix}$

6. $\begin{bmatrix} -3 & 0 \\ 0 & -2 \end{bmatrix}$

7. $\begin{bmatrix} 2 & -1 \\ 2 & 2 \end{bmatrix}$

8. $\begin{bmatrix} 4 & 6 \\ 0 & 4 \end{bmatrix}$

9. $\begin{bmatrix} 3 & -3 \\ 0 & 0 \\ 1 & 1 \end{bmatrix}$

10. $\begin{bmatrix} 7 & 1 \\ 5 & 5 \\ 0 & 0 \end{bmatrix}$

11. $\begin{bmatrix} -3 & 1 \\ 6 & -2 \\ 6 & -2 \end{bmatrix}$

12. $\begin{bmatrix} 1 & 1 \\ 0 & 1 \\ -1 & 1 \end{bmatrix}$

13. Find the SVD of $A = \begin{bmatrix} 3 & 2 & 2 \\ 2 & 3 & -2 \end{bmatrix}$ [Hint: Work with A^T .]

14. In Exercise 7, find a unit vector \mathbf{x} at which $A\mathbf{x}$ has maximum length.

15. Suppose the factorization below is an SVD of a matrix A , with the entries in U and V rounded to two decimal places.

$$A = \begin{bmatrix} .40 & -.78 & .47 \\ .37 & -.33 & -.87 \\ -.84 & -.52 & -.16 \end{bmatrix} \begin{bmatrix} 7.10 & 0 & 0 \\ 0 & 3.10 & 0 \\ 0 & 0 & 0 \end{bmatrix} \times \begin{bmatrix} .30 & -.51 & -.81 \\ .76 & .64 & -.12 \\ .58 & -.58 & .58 \end{bmatrix}$$

- a. What is the rank of A^T ?
- b. Use this decomposition of A , with no calculations, to write a basis for $\text{Col } A$ and a basis for $\text{Nul } A$. [Hint: First write the columns of V .]

16. Repeat Exercise 15 for the following SVD of a 3×4 matrix A :

$$A = \begin{bmatrix} -.86 & -.11 & -.50 \\ .31 & .68 & -.67 \\ .41 & -.73 & -.55 \end{bmatrix} \begin{bmatrix} 12.48 & 0 & 0 & 0 \\ 0 & 6.34 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \times \begin{bmatrix} .66 & -.03 & -.35 & .66 \\ -.13 & -.90 & -.39 & -.13 \\ .65 & .08 & -.16 & -.73 \\ -.34 & .42 & -.84 & -.08 \end{bmatrix}$$

In Exercises 17–24, A is an $m \times n$ matrix with a singular value decomposition $A = U\Sigma V^T$, where U is an $m \times m$ orthogonal matrix, Σ is an $m \times n$ “diagonal” matrix with r positive entries and no negative entries, and V is an $n \times n$ orthogonal matrix. Justify each answer.

- 17. Show that if A is square, then $|\det A|$ is the product of the singular values of A .
- 18. Suppose A is square and invertible. Find a singular value decomposition of A^{-1} .
- 19. Show that the columns of V are eigenvectors of $A^T A$, the columns of U are eigenvectors of AA^T , and the diagonal

426 CHAPTER 7 Symmetric Matrices and Quadratic Forms

entries of Σ are the singular values of A . [Hint: Use the SVD to compute $A^T A$ and AA^T .]

- 20. Show that if P is an orthogonal $m \times m$ matrix, then PA has the same singular values as A .
- 21. Justify the statement in Example 2 that the second singular value of a matrix A is the maximum of $\|Ax\|$ as \mathbf{x} varies over all unit vectors orthogonal to \mathbf{v}_1 , with \mathbf{v}_1 a right singular vector corresponding to the first singular value of A . [Hint: Use Theorem 7 in Section 7.3.]
- 22. Show that if A is an $n \times n$ positive definite matrix, then an orthogonal diagonalization $A = PDP^T$ is a singular value decomposition of A .
- 23. Let $U = [\mathbf{u}_1 \cdots \mathbf{u}_m]$ and $V = [\mathbf{v}_1 \cdots \mathbf{v}_n]$, where the \mathbf{u}_i and \mathbf{v}_i are as in Theorem 10. Show that $A = \sigma_1 \mathbf{u}_1 \mathbf{v}_1^T + \sigma_2 \mathbf{u}_2 \mathbf{v}_2^T + \cdots + \sigma_r \mathbf{u}_r \mathbf{v}_r^T$.
- 24. Using the notation of Exercise 23, show that $A^T \mathbf{u}_j = \sigma_j \mathbf{v}_j$ for $1 \leq j \leq r = \text{rank } A$.
- 25. Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. Describe how to find a basis \mathcal{B} for \mathbb{R}^n and a basis \mathcal{C} for \mathbb{R}^m such that the

matrix for T relative to \mathcal{B} and \mathcal{C} is an $m \times n$ “diagonal” matrix.

[M] Compute an SVD of each matrix in Exercises 26 and 27. Report the final matrix entries accurate to two decimal places. Use the method of Examples 3 and 4.

26. $A = \begin{bmatrix} -18 & 13 & -4 & 4 \\ 2 & 19 & -4 & 12 \\ -14 & 11 & -12 & 8 \\ -2 & 21 & 4 & 8 \end{bmatrix}$

27. $A = \begin{bmatrix} 6 & -8 & -4 & 5 & -4 \\ 2 & 7 & -5 & -6 & 4 \\ 0 & -1 & -8 & 2 & 2 \\ -1 & -2 & 4 & 4 & -8 \end{bmatrix}$

28. [M] Compute the singular values of the 4×4 matrix in Exercise 9 in Section 2.3, and compute the condition number σ_1/σ_4 .

29. [M] Compute the singular values of the 5×5 matrix in Exercise 10 in Section 2.3, and compute the condition number σ_1/σ_5 .