

MATH 2551 J/HP Midterm 2
VERSION A
Spring 2026
COVERS SECTIONS 12.1-12.5, 13.1-13.4, 14.1-14.2

Full name: _____ GT ID: _____

Honor code statement: I will abide strictly by the Georgia Tech honor code at all times. I will not use a calculator. **I do not have a phone within reach**, and I will not reference any website, application, or other CAS-enabled service. I will not consult with my notes or anyone during this exam. I will not provide aid to anyone else during this exam.

() All of the knowledge presented in this exam is entirely my own. I am initialing to the left to attest to my integrity.

Read all instructions carefully before beginning.

- Print your name and GT ID neatly above.
- You have 75 minutes to take the exam.
- You may not use aids of any kind.
- Please show your work [J] and annotate your work using proper notation [N].
- Good luck!

Question	Points
1	2
2	2
3	2
4	5
5	5
6	4
7	6
8	8
9	8
10	8
Total:	50

For T/F problems choose whether each statement is true or false. If the statement is *always* true, pick true. If the statement is *ever* false, pick false. For all problems on this page please be sure to neatly fill in the bubble corresponding to your answer choice. [A]

1. (2 points) If $f(x, y)$ is differentiable at the point $(2, 1)$, then the vector

$$\langle f_x(2, 1), f_y(2, 1), -1 \rangle$$

is normal to the plane tangent to the graph of $z = f(x, y)$ at the point $(2, 1)$.

☐ **TRUE**


☐ **FALSE**

2. (2 points) Suppose $w = f(x, y)$ is a differentiable function and $x(t), y(t)$ are both differentiable. Give an equation for the formula for $\frac{dw}{dt}$. [AN]

3. (2 points) Suppose S is a surface $F(x, y, z) = 7$ and $P(2, -1, 3)$ is a point on S . If $\nabla F(P) = \langle 4, 5, -6 \rangle$ then what is the equation for the tangent plane of S at $P(2, -1, 3)$? [AN]


4. (5 points) Sketch R the region of integration and switch the order of integration. That is, rewrite the integral in the order $dx dy$. *Do not evaluate!* [AN]

$$\int_0^2 \int_0^{x^2} \cos(x^3) dy dx$$




5. (5 points) Sketch R the region of integration and set up an integral in polar coordinates for evaluating $\iint_R f(x, y) dA$. Note that R is the region inside the circle of radius $r = 2$ which is over the line $y = x$. *Do not evaluate!* [AN]

$$R = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 \leq 4, y \geq x\}, \text{ and } f(x, y) = y.$$




Volume=



6. (4 points) In this problem, you will work with the function $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ whose contour plot near the origin is shown below. [AN]

- (a) Determine the sign $(+, -, 0)$ of the directional derivative at the point S in the direction $\langle 0, -2 \rangle$.
- (b) Determine the sign $(+, -, 0)$ of the directional derivative at the point Q in the direction $\langle 1, 0 \rangle$.
- (c) Draw a non-zero vector $\vec{u}$ at the point P for which $D_{\vec{u}}f(P) = 0$ the directional derivative of f in the direction of $\vec{u}$ equals 0 at the point P .



7. (6 points) Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ be the function $f(x, y) = e^{xy}$ and $\mathbf{r}(s, t) : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the function

$$\mathbf{r}(s, t) = \begin{bmatrix} s/t \\ s + t \end{bmatrix}.$$



- (a) Find Df .
- (b) Find $D\mathbf{r}$.
- (c) Evaluate $\mathbf{r}(2, 1)$ and $D\mathbf{r}|_{(s,t)=(2,1)}$.
- (d) Finally, use the Chain Rule to evaluate $D(f(\mathbf{r}(s, t)))|_{(s,t)=(2,1)}$.
Please put the answer to part (d) in the box below.

[AJN]

(d)

8. (8 points) Sketch the region R including labels for the axes, and labels for each boundary curve and each intersection point between curves. Then, compute the average value of $f(x, y) = y$ over the region R which is the triangle with vertices $(-2, 0)$, $(2, 0)$, and $(1, 3)$.

[AJN]



9. (8 points) Find all critical points and use the Hessian matrix to classify the critical points of the function. [AJN]

$$f(x, y) = x^2 + xy + 3x + 2y + 5$$

-
10. (8 points) Use the method of Lagrange Multipliers to find the maximum and minimum value of $f(x, y) = x + 2y$ subject to the constraint $x^2 + y^2 = 5$. [AJN]

FORMULA SHEET

- Total Derivative: For $\mathbf{f}(x_1, \dots, x_n) = \langle f_1(x_1, \dots, x_n), \dots, f_m(x_1, \dots, x_n) \rangle$

$$D\mathbf{f} = \begin{bmatrix} (f_1)_{x_1} & (f_1)_{x_2} & \cdots & (f_1)_{x_n} \\ (f_2)_{x_1} & (f_2)_{x_2} & \cdots & (f_2)_{x_n} \\ \vdots & \ddots & \cdots & \vdots \\ (f_m)_{x_1} & (f_m)_{x_2} & \cdots & (f_m)_{x_n} \end{bmatrix}$$

- Linearization: Near $\mathbf{a}$, $f(\mathbf{x}) \approx L(\mathbf{x}) = f(\mathbf{a}) + Df(\mathbf{a})(\mathbf{x} - \mathbf{a})$
- Chain Rule: If $h = g(f(\mathbf{x}))$ then $Dh(\mathbf{x}) = Dg(f(\mathbf{x}))Df(\mathbf{x})$
- Implicit Differentiation: If z is implicitly given in terms of x and y by $F(x, y, z) = c$, then $\frac{\partial z}{\partial x} = \frac{-F_x}{F_z}$ and $\frac{\partial z}{\partial y} = \frac{-F_y}{F_z}$.
- Directional Derivative: If $\mathbf{u}$ is a unit vector, $D_{\mathbf{u}}f(P) = Df(P)\mathbf{u} = \nabla f(P) \cdot \mathbf{u}$
- The tangent line to a level curve of $f(x, y)$ at (a, b) is $0 = \nabla f(a, b) \cdot \langle x - a, y - b \rangle$
- The tangent plane to a level surface of $f(x, y, z)$ at (a, b, c) is

$$0 = \nabla f(a, b, c) \cdot \langle x - a, y - b, z - c \rangle.$$

- Hessian Matrix: For $f(x, y)$, $Hf(x, y) = \begin{bmatrix} f_{xx} & f_{yx} \\ f_{xy} & f_{yy} \end{bmatrix}$
- Second Derivative Test: If (a, b) is a critical point of $f(x, y)$ then
 1. If $\det(Hf(a, b)) > 0$ and $f_{xx}(a, b) < 0$ then f has a local maximum at (a, b)
 2. If $\det(Hf(a, b)) > 0$ and $f_{xx}(a, b) > 0$ then f has a local minimum at (a, b)
 3. If $\det(Hf(a, b)) < 0$ then f has a saddle point at (a, b)
 4. If $\det(Hf(a, b)) = 0$ the test is inconclusive

- Area/volume: $\text{area}(R) = \iint_R dA$

- Trig identities: $\sin^2(x) = \frac{1}{2}(1 - \cos(2x))$, $\cos^2(x) = \frac{1}{2}(1 + \cos(2x))$

- Average value: $f_{\text{avg}} = \frac{\iint_R f(x, y) dA}{\text{area of } R}$

- Polar coordinates: $x = r \cos(\theta)$, $y = r \sin(\theta)$, $dA = r \, dr \, d\theta$

SCRATCH PAPER - PAGE LEFT BLANK