

Math 1553, Quiz 4 (2.5–3.1), Fall 2026, Version A

Name (ALL CAPS)		GT ID		Section	
----------------------------------	--	--------------	--	----------------	--

Please write your name in ALL CAPS. For questions with bubbles, either fill in the bubble completely or leave it blank. **Do not** mark any bubble with “X” or “/” or any such intermediate marking. Anything other than a blank or filled bubble will be given a score of zero.

1. (1 point) If $Ax = b$ is a consistent system of linear equations for some vector b in $\mathbf{R}^m$ and $T(x) = Ax$, then b is in the range of T .

☐ TRUE ☐ FALSE

2. (1 point) Which of the following sets of vectors are linearly independent? Fill in the bubble for all that apply.

☐ $\left\{ \begin{pmatrix} 1 \\ 2 \end{pmatrix}, \begin{pmatrix} 2 \\ 1 \end{pmatrix} \right\}$

 ☐ $\left\{ \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix}, \begin{pmatrix} 2 \\ 3 \\ 1 \end{pmatrix}, \begin{pmatrix} 5 \\ 6 \\ 7 \end{pmatrix} \right\}$

3. (1 point) Let S be the set of all vectors $\begin{pmatrix} x \\ y \end{pmatrix}$ in $\mathbf{R}^2$ satisfying $|x| = |y|$. Which of the properties of a subspace of $\mathbf{R}^2$ does S satisfy? Answer yes or no in each case.

(i) Is S closed under scalar multiplication? In other words, if u is in S and c is a scalar, is cu in S ? ☐ YES ☐ NO

(ii) Is S closed under addition? In other words, if u and v are in S , must $u + v$ also be in S ? ☐ YES ☐ NO